



NCCN Clinical Practice Guidelines in Oncology™

Rectal Cancer

V.1.2009

Continue

www.nccn.org

These guidelines are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network makes no representations nor warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way. These guidelines are copyrighted by National Comprehensive Cancer Network. All rights reserved. These guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2009.

National Comprehensive Cancer Network, Inc.

End User License Agreement for the NCCN Clinical Practice Guidelines in Oncology

IMPORTANT READ CAREFULLY: This End User License Agreement (the “Agreement”) is a legal agreement between you (either an individual or an entity) and the National Comprehensive Cancer Network, Inc. (“NCCN”) covering your use of one (1) PDF version of the NCCN Clinical Practice Guidelines in Oncology, which, together with any updates thereto, are hereinafter referred to collectively as the “Guidelines.” BY VIEWING OR OTHERWISE ACCESSING THE DATA CONTAINED IN THIS PDF, YOU AGREE TO BE BOUND BY THE TERMS AND CONDITIONS OF THIS LICENSE AGREEMENT. IF YOU DO NOT AGREE WITH THE TERMS OF THIS LICENSE AGREEMENT, DO NOT VIEW, ACCESS, OR USE THIS PDF AND RETURN THE PDF TO NCCN.

1. **Grant of License.** NCCN hereby grants to you a non-transferable, non-exclusive, limited license to access and use the Guidelines subject to the terms set forth in this License Agreement.

2. **Proprietary Rights.** You acknowledge that, as between you and NCCN, NCCN is the owner of all right, title and interest in and to the Guidelines, including, without limitation, all copyrights, trademarks, goodwill, derivative works, and other intellectual property and proprietary rights related thereto. Except for the limited rights expressly enumerated herein, you are not granted any rights relating to copyrights, trade names, trademarks (whether registered or unregistered) or any other rights, franchises or licenses with respect to the Guidelines or NCCN. You hereby agree that you shall not at any time dispute, challenge, or contest, directly or indirectly, NCCN's right, title and interest in and to the Guidelines, or assist or aid others to do so.

3. **Restrictions on Use.** You may not copy, transfer, reproduce, or create derivative works from, any part of the Guidelines for any reason. You may make and retain file copies of “Insubstantial Portions” of the Guidelines solely for your internal purposes. “Insubstantial Portions” means a quantity of data from the Guidelines that would not reasonably substitute for a comprehensive copy of the Guidelines and would not prejudice or diminish NCCN's advantage in licensing the Guidelines for commercial gain. Notwithstanding the foregoing, you may include Insubstantial Portions of the Guidelines in materials prepared in the ordinary course of your business for re-distribution to patients in connection with the delivery of your principal services. Any such materials shall cite NCCN as the source of the Guidelines and provide notice of NCCN's copyrights and other proprietary rights, as follows: © National Comprehensive Cancer Network, Inc. 2001-2009. NCCN and NATIONAL COMPREHENSIVE CANCER NETWORK are registered trademarks of National Comprehensive Cancer Network, Inc. You shall provide NCCN with examples of re-distributed materials including any portion of the Guidelines upon NCCN's reasonable request, but shall not be required to provide confidential information to NCCN. You agree to immediately cease any such re-distribution on receipt of notice from NCCN that, in NCCN's reasonable judgment, such re-distribution involves more than an Insubstantial Portion of the Guidelines or is otherwise in violation of this Agreement.

4. **Limited Warranty; Disclaimers; Limitation of Damages.** The Guidelines are a statement of consensus of its authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment.

NCCN MAKES NO WARRANTIES CONCERNING THE GUIDELINES OR ANY ELECTRONIC DELIVERY MEDIA SUPPLIED BY NCCN, WHICH ARE PROVIDED “AS IS.” NCCN DISCLAIMS ALL WARRANTIES, EXPRESS OR IMPLIED INCLUDING, WITHOUT LIMITATION, THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. NCCN DOES NOT WARRANT THE ACCURACY OR COMPLETENESS OF THE GUIDELINES AND DOES NOT WARRANT OR GUARANTEE OR MAKE ANY REPRESENTATION REGARDING THE USE OR THE RESULTS OF THE USE OF THE GUIDELINES. IN NO EVENT SHALL NCCN OR ITS MEMBERS BE LIABLE FOR ANY INCIDENTAL, INDIRECT, SPECIAL, PUNITIVE OR CONSEQUENTIAL DAMAGES ARISING OUT OF OR IN CONNECTION WITH THE LICENSE GRANTED UNDER THIS AGREEMENT OR USE OF THE GUIDELINES INCLUDING, WITHOUT LIMITATION, LOSS OF LIFE, LOSS OF DATA, LOSS OF INCOME OR PROFIT, OR OTHER LOSSES SUSTAINED AS A RESULT OF INJURY TO ANY PERSON, OR LOSS OR DAMAGE TO PROPERTY, OR CLAIMS OF THIRD PARTIES, EVEN IF NCCN HAS BEEN ADVISED OF THE POSSIBILITY OF SUCH DAMAGES.

SOME JURISDICTIONS DO NOT ALLOW THE LIMITATION OF IMPLIED WARRANTIES OR LIABILITY FOR INCIDENTAL, INDIRECT, SPECIAL, PUNITIVE OR CONSEQUENTIAL DAMAGES, SO THE ABOVE LIMITATIONS MAY NOT APPLY.

FOR ANY CLAIM YOU MAY HAVE AGAINST NCCN UNDER THIS AGREEMENT, YOUR EXCLUSIVE REMEDY AND NCCN'S ENTIRE LIABILITY SHALL BE TO PROVIDE REPLACEMENT GUIDELINES TO YOU.

5. **Trademarks.** NCCN and the NATIONAL COMPREHENSIVE CANCER NETWORK are trademarks (the “Marks”) of the National Comprehensive Cancer Network, Inc. and nothing in this Agreement shall constitute a license with respect to such trademarks. You shall not use the Marks or any confusingly similar Marks for any purpose, including, without limitation, for purposes of marketing or promoting your services, without the prior written approval of NCCN, which approval may be withheld in NCCN's sole discretion. Each approved use of the Marks shall require the independent written approval of NCCN.

6. **General.** This Agreement contains the entire agreement between NCCN and you relating to its subject matter. No amendment, change, or modification of this Agreement shall be binding on either party unless mutually agreed to by the parties in writing. If any provision of this Agreement is held to be invalid or unenforceable by a court of competent jurisdiction, the validity and enforceability of the remaining provisions shall not be affected thereby. This Agreement will be governed by and construed in accordance with the laws of the Commonwealth of Pennsylvania without giving any effect to the conflict of law provisions thereof, and each party agrees to submit to personal jurisdiction in the federal and state courts of Pennsylvania and waives any objection to venues in said courts. This Agreement will not be governed by the United Nations Conventions of Contracts for the International Sale of goods, the application of which is expressly excluded. You agree that the Guidelines will not be shipped, transferred or exported into any country or used in any manner prohibited by the United States Export Administration Act, or any other export laws, restrictions. This Agreement will terminate automatically upon failure by you to comply with its terms.

BY ACCESSING THE DATA CONTAINED IN THIS PDF, YOU ACKNOWLEDGE THAT YOU HAVE READ THIS AGREEMENT, UNDERSTAND IT, AND AGREE TO BE BOUND BY ITS TERMS AND CONDITIONS.

[Click Here to Continue](#)

NCCN Rectal Cancer Panel Members

*Paul F. Engstrom, MD/Chair †
Fox Chase Cancer Center

Juan Pablo Arnoletti, MD ¶
University of Alabama at Birmingham
Comprehensive Cancer Center

*Al B. Benson, III, MD †
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Yi-Jen Chen, MD, PhD §
City of Hope

Michael A. Choti, MD ¶
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Harry S. Cooper, MD ≠
Fox Chase Cancer Center

Anne Covey, MD ϕ
Memorial Sloan-Kettering Cancer Center

Raza A. Dilawari, MD ¶
St. Jude Children's Research
Hospital/University of Tennessee Cancer
Institute

Dayna S. Early, MD ≠
Siteman Cancer Center at Barnes-Jewish
Hospital and Washington University
School of Medicine

Peter C. Enzinger, MD †
Dana-Farber/Brigham and Women's
Cancer Center

Marwan G. Fakih, MD †
Roswell Park Cancer Institute

James Fleshman, Jr., MD ¶
Siteman Cancer Center at Barnes-Jewish
Hospital and Washington University School
of Medicine

Charles Fuchs, MD †
Dana-Farber/Brigham and Women's Cancer
Center | Massachusetts General Hospital
Cancer Center

Jean L. Grem, MD †
UNMC Eppley Cancer Center at The
Nebraska Medical Center

Krystyna Kiel, MD §
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

James A. Knol, MD ¶
University of Michigan Comprehensive
Cancer Center

Lucille A. Leong, MD †
City of Hope Cancer Center

Edward Lin, MD †
Fred Hutchinson Cancer Research
Center/Seattle Cancer Care Alliance

Mary F. Mulcahy, MD ‡
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Sujata Rao, MD †
Fred Hutchinson Cancer Research
Center/Seattle Cancer Care Alliance

David P. Ryan, MD ≠
Massachusetts General Hospital Cancer Center

*Leonard Saltz, MD † ‡ P
Memorial Sloan-Kettering Cancer Center

David Shibata, MD ¶
H. Lee Moffitt Cancer Center and Research
Institute at the University of South Florida

John M. Skibber, MD ¶
The University of Texas M. D. Anderson Cancer
Center

Constantinos Sofocleous, MD, PhD ϕ
Memorial Sloan-Kettering Cancer Center

James Thomas, MD
Arthur G. James Cancer Hospital & Richard J.
Solove Research Institute at The Ohio State
University

Alan P. Venook, MD † ‡
UCSF Comprehensive Cancer Center

Christopher Willett, MD §
Duke Comprehensive Cancer Center

† Medical Oncology
§ Radiotherapy/Radiation oncology
¶ Surgery/Surgical oncology
≠ Pathology
‡ Hematology/Hematology Oncology
P Internal medicine
≠ Gastroenterology
ϕ Diagnostic/Interventional Radiology
* Writing Committee Member

Continue

[NCCN Guidelines Panel Disclosures](#)

Table of Contents

[NCCN Rectal Cancer Panel Members](#)

[Summary of the Guidelines Updates](#)

Clinical Presentations and Primary Treatment:

- [Pedunculated polyp with invasive cancer \(REC-1\)](#)
- [Sessile polyp with invasive cancer \(REC-1\)](#)
- [Rectal cancer appropriate for resection \(REC-2\)](#)
 - ▶ [T1-2, N0: Primary and Adjuvant Treatment \(REC-3\)](#)
 - ▶ [T3, N0 or T any, N1-2: Primary and Adjuvant Treatment \(REC-4\)](#)
 - ▶ [T4 and/or locally unresectable: Primary and Adjuvant Treatment \(REC-4\)](#)
 - ▶ [T any, N any, M1: Resectable Metastases Treatment and Surveillance \(REC-5\)](#)
 - ▶ [T any, N any, M1: Unresectable Metastases or Medically Inoperable Treatment \(REC-6\)](#)

[Surveillance \(REC-7\)](#)

[Recurrence and Workup \(REC-8\)](#)

[Postoperative CEA Elevation \(REC-8\)](#)

[Principles of Pathologic Review \(REC-A\)](#)

[Principles of Surgery \(REC-B\)](#)

[Principles of Adjuvant Therapy \(REC-C\)](#)

[Principles of Radiation Therapy \(REC-D\)](#)

[Chemotherapy for Advanced or Metastatic Disease \(REC-E\)](#)

[Principles of Survivorship \(REC-F\)](#)

[Print the Rectal Cancer Guideline](#)

These guidelines are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network makes no representations nor warranties of any kind whatsoever regarding their content, use, or application and disclaims any responsibility for their application or use in any way. These guidelines are copyrighted by National Comprehensive Cancer Network. All rights reserved. These guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2009.

[For help using these documents or for more information about the NCCN Guidelines and the Complete Library of Clinical Practice Guidelines in Oncology, please click here](#)

[Staging](#)

[Discussion](#)

[References](#)

This manuscript is being updated to correspond with the newly updated algorithm.

Clinical Trials: The NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN member institutions, [click here: nccn.org/clinical_trials/physician.html](#)

NCCN Categories of Evidence and Consensus: All recommendations are Category 2A unless otherwise specified.

See [NCCN Categories of Evidence and Consensus](#)

Summary of the Guidelines updates

Summary of changes in the 1.2009 version of the Rectal Cancer Guidelines from the 3.2008 version include:

[REC-2](#)

- “Rigid” was added to proctoscopy in the workup section.

[REC-5](#)

- The following therapy option was added for patients with synchronous resectable metastases: Combination chemotherapy for 2-3 months, followed by chemotherapy/RT, followed by staged or synchronous resection.

[REC-7](#)

- A link was added to Principles of Survivorship in the Surveillance section.

[REC-8](#)

- For isolated, pelvic/anastomotic recurrence, RT was clarified as IORT if given during resection.

[REC-9](#)

- Unresectable disease was defined as including “potentially convertible” and “unconvertible”. Further guidance and a description of these categories was added to the Principles of Surgery section (REC-B 2 of 3).
- The recommendation for “re-evaluation for conversion to resectable every 2 mo” was added after primary treatment.
- Footnote “z” was added recommending that patients should be evaluated by a multidisciplinary team including surgical consultation for potentially resectable patients.
- The recommendation for hepatic artery infusion was moved from the body of the algorithm to footnote “aa”.
- The treatment option of observation was moved from the footnote into the body of the algorithm after primary treatment.
- There is a new footnote “bb” specifying that therapy should be considered for a maximum of 6 months.

[REC-10](#)

- Footnote “z” was added recommending that patients should be evaluated by a multidisciplinary team including surgical consultation for potentially resectable patients.
- The clarification of “2-3 months” was added for neoadjuvant chemotherapy.
- The recommendation for hepatic artery infusion was moved from the body of the algorithm to footnote “aa”.
- The treatment option of observation was moved from the footnote into the body of the algorithm after primary treatment.
- There is a new footnote “bb” specifying that therapy should be considered for a maximum of 6 months.

[Continued](#)

Summary of the Guidelines updates

Summary of changes in the 1.2009 version of the Rectal Cancer Guidelines from the 3.2008 version include:

REC-A 3 of 4

- The KRAS Mutation testing section was added to provide further definition and direction for testing and use of results.
- The Evaluation of Mesorectum (TME) section was added

REC-A 4 of 4

- References 37-39 were added to support KRAS information. References 40-42 were added to support TME information.

REC-B 2 of 3

Liver - the following bullets were added to the page:

- Patients with resectable metastatic disease and primary tumor in place should have both sites resected with curative intent. These can be resected in one operation or as a staged approach, depending on the complexity of the hepatectomy or colectomy, comorbid diseases, surgical exposure, and surgeon expertise.
- When hepatic metastatic disease is not optimally resectable based on insufficient remnant liver volume, approaches utilizing preoperative portal vein embolization or staged liver resection can be considered.
- Some institutions use intra-arterial embolization in select patients with chemotherapy resistant/refractory disease, without obvious systemic disease, with predominant hepatic metastases (category 3).
- Conformal external beam radiation therapy should not be used unless the patient is symptomatic or in the setting of a clinical trial.

Lung - the following bullets were added to the page:

- Ablative techniques can be considered when unresectable and amenable to complete ablation.
- Patients with resectable synchronous metastases can be resected synchronously or using a staged approach.

NEW SECTION - There is a new section with recommendations for evaluating a patient for conversion to resectable disease.

REC-D

The following bullet was modified:

- Intensity modulated radiotherapy (IMRT) or tomotherapy should only be used in the setting of a clinical trial.

REC-E 1 of 6

- Patients appropriate for therapy - the following options were added for initial therapy: FOLFOX or FOLFIRI or CapeOX ± cetuximab (KRAS wild-type gene only), FOLFOXIRI with a category 2B designation.
- 5FU/leucovorin + bevacizumab was added as a treatment option for patients progressing after FOLFOXIRI. If patients progress on 5FU/leucovorin + bevacizumab, the recommended therapy options are cetuximab or panitumumab.

REC-E 2 of 6


- Cetuximab was added as a treatment option for patients not appropriate for intensive therapy with a category 2B designation.

REC-E 3 of 6

- Footnote 5 is new to the page: Combination therapy involving more than one biologic agent is not recommended.
- Footnote 10 is new to the page: Data are not mature for the addition of biologic agents to FOLFOXIRI.

REC-F

- Principles of Survivorship is a new section to the Guidelines.



^aAll patients with colon cancer should be counseled for family history. Patients with suspected hereditary non-polyposis colon cancer (HNPCC), familial adenomatous polyposis (FAP) and attenuated FAP, see the [NCCN Colorectal Cancer Screening Guidelines](#).

^bConfirm the presence of invasive cancer (pT1). pT1s has no biological potential to metastasize.

^cIt has not been established if molecular markers are useful in treatment determination (predictive markers) and prognosis. College of American Pathologists Consensus Statement 1999. Prognostic factors in colorectal cancer. Arch Pathol Lab Med 2000;124:979-994.

^d[See Principles of Pathologic Review \(REC-A\)](#) - Endoscopically removed malignant polyp.

[Back to Other Clinical Presentations \(Table of Contents\)](#)


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CLINICAL
PRESENTATION

WORKUP


CLINICAL STAGE



^aAll patients with colon cancer should be counseled for family history. Patients with suspected hereditary non-polyposis colon cancer (HNPCC), familial adenomatous polyposis (FAP) and attenuated FAP, see the [NCCN Colorectal Cancer Screening Guidelines](#).

^eT1-2, N0 should be based on assessment of endorectal ultrasound or MRI.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



^eT1-2, N0 should be based on assessment of endorectal ultrasound or MRI.

^fSee Principles of Surgery (REC-B).

^gHigh risk features include positive margins, lymphovascular invasion and poorly differentiated tumors.

^hSee Principles of Adjuvant Therapy (REC-C).


ⁱSee Principles of Radiation Therapy (REC-D).

^jThe use of FOLFOX or capecitabine is an extrapolation from the available data in colon cancer. Trials are still pending in rectal cancer.

^kData regarding the use of capecitabine/RT is limited and no phase III randomized data are available. Trials are pending. Kim J-Sang, Kim J-Sung, Cho, M, et al Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. Int J Radiation Oncology Biol Phys 2002;54(2):403-408.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



^fSee Principles of Surgery (REC-B).

^hSee Principles of Adjuvant Therapy (REC-C).

ⁱSee Principles of Radiation Therapy (REC-D).

^jThe use of FOLFOX or capecitabine is an extrapolation from the available data in colon cancer. Trials are still pending in rectal cancer.

^kData regarding the use of capecitabine/RT is limited and no phase III randomized data are available. Trials are pending. Kim J-Sang, Kim J-Sung, Cho, M, et al. Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. Int J Radiation Oncology Biol Phys 2002;54(2):403-408.

^lThe use of agents other than fluoropyrimidines are not recommended concurrently with RT.


^mFor patients with proximal T3, N0 disease with clear margins and favorable prognostic features, the incremental benefit of RT is likely to be small. Consider chemotherapy alone.

ⁿPostoperative therapy is indicated in all patients who receive preoperative therapy, regardless of the surgical pathology results.

^oAn ongoing Intergroup trial compares 5-FU/leucovorin, FOLFOX, and FOLFIRI after surgery.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



^fSee Principles of Surgery (REC-B).

^hSee Principles of Adjuvant Therapy (REC-C).

ⁱSee Principles of Radiation Therapy (REC-D).

^jThe use of FOLFOX or capecitabine is an extrapolation from the available data in colon cancer. Trials are still pending in rectal cancer.

^kData regarding the use of capecitabine/RT is limited and no phase III randomized data are available. Trials are pending. Kim J-Sang, Kim J-Sung, Cho, M et al Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. Int J Radiation Oncology Biol Phys 2002;54(2):403-408.

^oAn ongoing Intergroup trial compares 5-FU/leucovorin, FOLFOX, and FOLFIRI after surgery.

^pDetermination of tumor KRAS gene status. See Principles of Pathologic Review (REC-A 3 of 4) - KRAS Mutation Testing.

^qThe safety of administering bevacizumab pre or postoperatively, in combination with 5-FU-based regimens, has not been adequately evaluated. There should be at least a 6 wk interval between the last dose of bevacizumab and elective surgery. There is an increased risk of stroke and other arterial events especially in age ≥ 65. The use of bevacizumab may interfere with wound healing.


^rRT only recommended for patients at relative risk for pelvic recurrence.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CLINICAL STAGE

PRIMARY TREATMENT



^kData regarding the use of capecitabine/RT is limited and no phase III randomized data are available. Trials are pending. Kim J-Sang, Kim J-Sung, Cho, M et al Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. Int J Radiation Oncology Biol Phys 2002;54(2):403-408.

^PDetermination of tumor KRAS gene status. [See Principles of Pathologic Review \(REC-A 3 of 4\)](#) - KRAS Mutation Testing.

^S[See Chemotherapy for Advanced or Metastatic Disease \(REC-E\)](#).

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

SURVEILLANCE

- History and physical every 3-6 mo for 2 y, then every 6 mo for a total of 5 y
- CEA[†] every 3-6 mo for 2 y, then every 6 mo for a total of 5 y for T2 or greater lesions
- Chest/abdominal/pelvic CT annually x 3 y for patients at high risk for recurrence^{u,v}
- Colonoscopy in 1 y except if no preoperative colonoscopy due to obstructing lesion, colonoscopy in 3-6 mo
 - ▶ If abnormal, repeat in 1 y
 - ▶ If no advanced adenoma,^w repeat in 3 y, then every 5 y^x
- Consider proctoscopy every 6 mo x 5 y for patients status post LAR^y
- PET scan is not routinely recommended
- See [Principles of Survivorship \(REC-F\)](#)



[†]If patient is a potential candidate for resection of isolated metastasis.

^uDesch CE, Benson III AB, Somerfield MR, et al. Colorectal cancer surveillance: 2005 update of the American Society of Clinical Oncology Practice Guideline. J Clin Oncol 2005;23(33):8512-8519.

^vCT scan may be useful for patients at high risk for recurrence (eg, lymphatic or venous invasion by tumor, or poorly differentiated tumors).


^wVillous polyp, polyp > 1 cm, or high grade dysplasia.

^xRex DK, Kahi CJ, Levin B, et al. Guidelines for colonoscopy surveillance after cancer resection: a consensus update by the American Cancer Society and the US Multi-Society Task Force on Colorectal Cancer. Gastroenterology 2006;130(6):1865-71.

^yPatients with rectal cancer should also undergo limited endoscopic evaluation of the rectal anastomosis to identify local recurrence. Optimal timing for surveillance is not known. No specific data clearly support rigid versus flexible proctoscopy. The utility of routine endoscopic ultrasound for early surveillance is not defined.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




ⁱSee Principles of Radiation Therapy (REC-D).

^PDetermination of tumor KRAS gene status. See Principles of Pathologic Review (REC-A 3 of 4) - KRAS Mutation Testing.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRIMARY TREATMENT



^f See Principles of Surgery (REC-B).

^PDetermination of tumor KRAS gene status. See Principles of Pathologic Review (REC-A 3 of 4) - KRAS Mutation Testing.


^ZPatients should be evaluated by a multidisciplinary team including surgical consultation for potentially resectable patients.

^{aa}Hepatic artery infusion ± systemic 5-FU/leucovorin (category 2B) is also an option at institutions with experience in both the surgical and medical oncologic aspects of this procedure.

^{bb}Therapy may be considered for a maximum of 6 months.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



^fSee Principles of Surgery (REC-B).

^zPatients should be evaluated by a multidisciplinary team including surgical consultation for potentially resectable patients.

^{aa}Hepatic artery infusion ± systemic 5-FU/leucovorin (category 2B) is also an option at institutions with experience in both the surgical and medical oncologic aspects of this procedure.

^{bb}Therapy may be considered for a maximum of 6 months.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF PATHOLOGIC REVIEW (1 of 4)

Endoscopically removed malignant polyps

- A malignant polyp is defined as one with cancer invading through the muscularis mucosae and into the submucosa (pT1). pT1S is not considered a “malignant polyp.”
- Favorable histological features grade 1 or 2, no angiolymphatic invasion and negative margin of resection. There is no consensus as to the definition of what constitutes a positive margin of resection. A positive margin has been defined as 1) tumor < 1 mm from the transected margin, 2) tumor < 2 mm from the transected margin, 3) tumor cells present within the diathermy of the transected margin.¹⁻⁴
- Unfavorable histological features grade 3 or 4, or angiolymphatic invasion, or a “positive margin.” See above for definition of a positive margin.
- There is controversy as to whether malignant colorectal polyps with a sessile configuration can be successfully treated by endoscopic removal. The literature seems to indicate that endoscopically removed sessile malignant polyps have a significantly greater incidence of adverse outcome (residual disease, recurrent disease, mortality, hematogenous metastasis, but not lymph node metastasis) than do polypoid malignant polyps. However, when one closely looks at the data, configuration by itself is not a significant variable for adverse outcome and endoscopically removed malignant sessile polyps with grade I or II histology, negative margin, and no lymphovascular invasion can be successfully treated with endoscopic polypectomy.³⁻⁷

Transanal excision

- Favorable histopathological features: < 3 cm size, T1 or T2 (use caution in T2 due to high recurrence rate [see REC-B](#)), grade I or II, no lymphatic or venous invasion, negative margins.^{8,9}
- Unfavorable histopathological features: > 3 cm in size, T1 or T2, with grade III, or lymphovascular invasion, or positive margin.⁸⁻¹⁰

Rectal cancer appropriate for resection

- Histological confirmation of primary malignant rectal neoplasm.

Pathological stage

- The following parameters should be reported.
 - ▶ Grade of the cancer
 - ▶ Depth of penetration, (T) the T stage is based on viable tumor. Acellular mucin pools are not considered residual tumor in those cases treated with neoadjuvant therapy.
 - ▶ Number of lymph nodes evaluated and number positive (N). Acellular mucin pools are not considered residual tumor in those cases treated with neoadjuvant therapy.
 - ▶ Status of proximal, distal, and circumferential (radial) margins.¹¹⁻¹²
 - ▶ A positive circumferential resection margin (CRM) has been defined as < 1 mm or < 2 mm depending on the publication¹³⁻¹⁴

[See Staging \(ST-1\)](#)

[See Lymph node evaluation and sentinel lymph node on page 2 of 4 REC-A](#)

[See KRAS Mutation Testing page 3 of 4 REC-A](#)

[See footnotes on page 4 of 4 REC-A](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF PATHOLOGIC REVIEW (2 of 4)

Lymph node evaluation

- The AJCC and College of American Pathologists recommend examination of a minimum of 12 lymph nodes to accurately identify stage II colorectal cancers.^{11,12,15} The literature lacks consensus as to what is the minimal number of lymph nodes to accurately identify stage II cancer. The minimal number of nodes has been reported as >7, >9, >13, >20, >30.¹⁶⁻²³ Most of these studies have combined rectal and colon cancers and reflect those cases with surgery as the initial treatment. Two studies confined only to rectal cancer have reported 14 and > 10 lymph nodes as the minimal number to accurately identify stage II rectal cancer.^{19,22} The number of lymph nodes retrieved can vary with age of the patient, gender, tumor grade and tumor site.¹⁶ For stage II (pN0) colon cancer, if less than 12 lymph nodes are initially identified, it is recommended that the pathologist go back to the specimen and resubmit more tissue of potential lymph nodes. If 12 lymph nodes are still not identified, a comment in the report should indicate that an extensive search for lymph nodes was undertaken. The mean number of lymph nodes retrieved from rectal cancers treated with neoadjuvant therapy is significantly less than those treated by surgery alone (13 vs 19, $p < 0.05$, 7 vs 10, $p < 0.001$).^{24,25} If 12 lymph nodes is considered the number needed to accurately stage, stage II tumors, then only 20% of cases treated with neoadjuvant therapy had adequate lymph node sampling.²⁵ To date the number of lymph nodes needed to accurately stage neoadjuvant treated cases is unknown. However, it is not known what is the clinical significance of this in the neoadjuvant setting as postoperative therapy is indicated in all patients who receive preoperative therapy, regardless of the surgical pathology results.

Sentinel lymph node and detection of micrometastasis by immunohistochemistry

- Examination of the sentinel lymph node allows an intense histological and/or immunohistochemical investigation to detect the presence of metastatic carcinoma. Studies in the literature have been reported using multiple H & E sections and/or immunohistochemistry (IHC) to detect cytokeratin positive cells. While studies to date seem promising, there is no uniformity in the definition of what constitutes "true metastatic carcinoma." Confusion arises when isolated tumor cells (ITC) have been considered micrometastatic disease in contraindication to true micrometastasis (tumor aggregates > 0.2 mm to < 2 mm in size). The significance of detection of single cells by IHC alone is controversial. Some studies have considered these to be micrometastasis, however, "consensus" recommends these to be considered ITC and not micrometastatic disease.²⁶⁻²⁸ While the 6th edition of the AJCC Cancer Staging²⁹ manual considers "tumor clusters" < 0.2 mm as isolated tumor cells (pN0) and not metastatic carcinoma, some have challenged this. Some investigators believe that size should not effect the diagnosis of metastatic cancer. They believe that tumor foci that show evidence of growth (eg, glandular differentiation, distension of sinus, or stromal reaction) should be diagnosed as a lymph node metastasis regardless of size.³⁰ Hermanek et al³¹ proposed isolated tumor cells to be defined as single tumor cells or small clusters (never more than a few cells clumped together) without evidence of extrasinusoidal stromal proliferation or reaction and no contact with or invasion of the vessel (lymphatic) wall.
- Some studies have shown that the detection of IHC cytokeratin positive cells in stage II (N0) colon cancer (defined by H & E) has a worse prognosis while others have failed to show this survival difference. In these studies, ITC were considered micrometastasis.³²⁻³⁶
- At the present time the use of sentinel lymph nodes and detection of cancer cells by IHC alone should be considered investigational and results used with caution in clinical management decisions.^{26-28,32-36}

[See Malignant polyp, rectal cancer appropriate for resection, and pathological stage on page 1 of 4 REC-A](#)

[See KRAS Mutation Testing page 3 of 4 REC-A](#)

[See footnotes on page 4 of 4 REC-A](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF PATHOLOGIC REVIEW (3 of 4)

KRAS Mutation Testing

- Mutations in codons 12 and 13 in exon 2 of the coding region of the KRAS gene predict lack of response to therapy with antibodies targeted to the epidermal growth factor receptor.^{37,38}
-
- Testing for Mutations in Codons 12 and 13 should be performed only in laboratories that are certified under the clinical laboratory improvement amendments of 1988 (CLIA – 88) as qualified to perform high complex clinical laboratory (molecular pathology) testing. No specific methodology is recommended (sequencing, hybridization, etc.).
-
- The testing can be performed on formalin fixed paraffin embedded tissue. The testing can be performed on the primary colorectal cancers and/or the metastasis as literature has shown that the KRAS mutations are similar in both specimen types.³⁹

Evaluation of Mesorectum (TME)

- The pathologist should evaluate the quality (completeness) of the mesorectum (only for low rectal cancer - distal 2/3).⁴⁰⁻⁴²

[See footnotes on page 4 of 4 REC-A](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF PATHOLOGIC REVIEW (3 of 3) - References

- ¹Volk EE, Goldblum JR, Petras RE, et al. Management and outcome of patients with invasive carcinoma arising in colorectal polyps. *Gastroenterology* 1995;109:1801-1807.
- ²Cooper HS, Deppisch LM, Gourley WK, et al. Endoscopically removed malignant colorectal polyps: clinical pathological correlations. *Gastroenterology* 1995;108:1657-1665.
- ³Ueno H, Mochizuki H, Hashiguchi Y, et al. Risk factors for an adverse outcome in early invasive colorectal carcinoma. *Gastroenterology* 2004;127:385-394.
- ⁴Seitz U, Bohnacker S, Seewald S, et al. Is endoscopic polypectomy an adequate therapy for malignant colorectal polyps? Presentation of 114 patients and review of the literature. *Dis Colon Rectum* 2004;47:1789-1797.
- ⁵Morson BC, Whiteway JE, Jones EA, et al. Histopathology and prognosis of malignant colorectal polyps treated by endoscopic polypectomy. *Gut* 1984;25:437-444.
- ⁶Haggitt RC, Glotzbach RE, Soffer EE, Wruble LD. Prognostic factors in colorectal carcinomas arising in adenomas: implications for lesions removed by endoscopic polypectomy. *Gastroenterology* 1985;89:328-336.
- ⁷Netzer P, Binck J, Hammer B, et al. Significance of histological criteria for the management of patients with malignant colorectal polyps. *Scand J Gastroenterol* 1997;323:915-916.
- ⁸Hager T, Gall FP, and Hermanek P. Local excision of cancer of the rectum. *Dis Colon Rect* 1983;26:149-151.
- ⁹Willett, CG, Tepper JE, Donnelly S, et al. Patterns of failure following local excision and local excision and postoperative radiation therapy for invasive rectal adenocarcinoma. *J Clin Oncol* 1989;7:1003-1008.
- ¹⁰Nascimbeni R, Burgart LJ, Nivatvongs S, and Larson DR. Risk of lymph node metastasis in T1 carcinoma of the colon and rectum. *Dis Colon Rectum* 2002;45:2001-2006.
- ¹¹Compton CC and Greene FL. The staging of colorectal cancer: 204 and beyond. *Cancer J Clin* 2004;54:295-308.
- ¹²Compton CC, Fielding LP, Burkhardt LJ, et al. Prognostic factors in colorectal cancer. College of American pathologists consensus statement. *Arch Pathol Lab Med* 2000;124:979-994.
- ¹³Nagtegaal ID, Merijnena M, Kranenbarg EK, et al. Circumferential margin involvement is still an important predictive local occurrence in rectal carcinoma. Not one millimeter but two millimeters is the limit. *Am J Surg Pathol* 2002;26:350-357.
- ¹⁴Wibe A, Rendedal PR, Svensson E, et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surgery* 2002;89 327-334.
- ¹⁵Sobin HL and Green EFL. TNM classification. Clarification of number of regional lymph node for PN0. *Cancer* 2001;92:452.
- ¹⁶Sarli L, Bader G, Lusco D, et al. Number of lymph nodes examined and prognosis of TNM stage II colorectal cancer. *European Journal of Cancer* 2005;41:272-279.
- ¹⁷Chaplin S, Scerottini G-P, Bosman FT, et al. For patients with Duke's B (TNM stage II) colorectal carcinoma, examination of six or fewer lymph nodes is related to poor prognosis. *Cancer* 1998;83:666-72.
- ¹⁸Maurel J, Launoy G, Grosclaude P, et al. Lymph node harvest reporting in patients with carcinoma of the large bowel. A French population-based study. *Cancer* 1998;82:1482-6.
- ¹⁹Pocard M, Panis Y, Malassagane B, et al. Assessing the effectiveness of mesorectal excision in rectal cancer. *Dis Colon Rectum* 1998;41:839-845.
- ²⁰Joseph NE, Sigurdson ER, Hamlin AL, et al. Accuracy of determining nodal negativity in colorectal cancer on the basis of number of nodes retrieved on resection. *Ann of Surg Oncol* 2003;10:213-218.
- ²¹Goldstein NS. Lymph node recurrences from 2427 PT3 colorectal resection specimens spanning 45 years. Recommendations for a minimum number of recovered lymph nodes based on predictive probabilities. *Am J Surg Pathol* 2002;26:179-189.
- ²²Tepper JE, O'Connell MJ, Niedzwiecki D, et al. Impact of number of nodes retrieved on outcome in patients with rectal cancer. *J Clin Oncol* 2001;19:157-162.
- ²³Scott KWM and Grace RH. Detection of lymph node metastasis and colorectal carcinoma before and after fat clearance. *Br J Surg* 1989;76:1165-1167.
- ²⁴Wichmann MW, Mollar C, Meyer G, et al. Effect of pre-operative radiochemotherapy on lymph node retrieval after resection of rectal cancer. *Arch Surg* 2002;137:206-210.
- ²⁵Baxter NN, Morris AM, Rothenberger DA, and Tepper JE. Impact of pre-operative radiation for rectal cancer on subsequent lymph node evaluation: population based analysis. *Int J Radiation Oncology Biol Phys* 2005;61:426-431.
- ²⁶Turner RR, Nora DT, Trochas D, and Bilchik AJ. Colorectal carcinoma in nodal staging. Frequency and nature of cytokeratin positive cells in sentinel and nonsentinel lymph nodes. *Arch Pathol Lab Med* 2003;127:673-679.
- ²⁷Wood TF, Nora DT, Morton DL, et al. One hundred consecutive cases of sentinel node mapping in early colorectal carcinoma. Detection of missed micrometastasis. *J Gastrointest Surg* 2002;6:322-330.
- ²⁸Wiese DA, Sha S, Badin J, et al. Pathological evaluation of sentinel lymph nodes in colorectal carcinoma. *Arch Pathol Lab Med* 2000;124:1759-1763.
- ²⁹AJCC Cancer Staging Manual, 6th ed. Greene FL, Page D, Balch C, et al (editors) Springer, New York, 2002:227.
- ³⁰Jass JB, O'Brien MJ, Riddell RH, Snover DC, on behalf of the Association of Directors of Anatomic and Surgical Pathology. Recommendations for the reporting of surgically resected specimens of colorectal carcinoma. *Hum Pathol* 2007;38:537-545.
- ³¹Hermanek P, Hutter RVP, Sobin LH, Wittekind CH. Classification of isolated tumor cells and micrometastasis. *Cancer* 1999;86:2668-73.
- ³²Noura S, Yamamoto H, Ohnishi T, et al. Comparative detection of lymph node micrometastasis of stage II colorectal cancer by reverse transcriptase polymerase chain reaction in immunohistochemistry. *J Clin Oncol* 2002;20:4232-4241.
- ³³Yasuda K, Adachi Y, Shiraiishi N, et al. Pattern of lymph node micrometastasis and prognosis of patients with colorectal cancer. *Ann Surg Oncol* 2001;8:300-304.
- ³⁴Noura S, Yamamoto H, Miyake Y, et al. Immunohistochemical assessment of localization of frequency of micrometastasis in lymph nodes of colorectal cancer. *Clin Cancer Research* 2002;8:759-767.
- ³⁵Oberg A, Stenling R, Tavelin B, Lindmark G. Are lymph node micrometastasis of any clinical significance in Duke stages A and B colorectal cancer? *Dis Colon Rectum* 1998;41:1244-1249.
- ³⁶Greenon JK, Isenhardt TCE, Rice R, et al. Identification of occult micrometastasis in pericolic lymph nodes of Duke's B colorectal cancer. Patient's using monoclonal antibodies against cytokeratin and CC49. Correlation with long term survival. *Cancer* 1994;73:563-9.
- ³⁷Lievre A, Bachatte J-B, Blige V, et al. KRAS mutations as an independent prognostic factor in patients with advanced colorectal cancer treated with Cetuximab. *J Clin Oncol* 2008;26:374-379.
- ³⁸Amado IG, Wolf M, Peters M, et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* 2008;26:1626-1634.
- ³⁹Etienne-Gimeldi M-C, Formenta J-L, Francoual M, et al. KRAS mutations in treatment outcome in colorectal cancer in patients receiving exclusive fluoropyrimidine. *Clin Cancer Research* 2008;14:4830-4835.
- ⁴⁰Parfitt JR and Driman KR. Total mesorectal excision specimen for rectal cancer: A review of its pathological assessment. *J Clin Pathol* 60:849-855, 2007.
- ⁴¹Jass JR, O'Brien MJ, Riddell RH, Snover DC. On behalf of the association of Directors of Anatomic and Surgical Pathology recommendations for the reporting of surgically resected specimens in colorectal carcinoma. *Human Pathol* 38:537-545, 2007.
- ⁴²Nagtegaal ID, Vandevelde CJA, Derworp EV, et al. Macroscopic evaluation of the rectal cancer resection margin: Clinical significance of the pathologist in quality control. *J Clin Oncol* 20: 1729-1734, 2002.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURGERY (1 of 3)

Transanal excision:**• Criteria**

- ▶ < 30% circumference of bowel
 - ▶ < 3 cm in size
 - ▶ Margin clear (> 3 mm)
 - ▶ Mobile, nonfixed
 - ▶ Within 8 cm of anal verge
 - ▶ T1 or T2 (use caution in T2, due to high recurrence rate)
 - ▶ Endoscopically removed polyp with cancer or indeterminate pathology
 - ▶ No lymphovascular (LVI) or perineural invasion
 - ▶ Well to moderately differentiated
 - ▶ No evidence of lymphadenopathy on pretreatment imaging
- When the lesion can be adequately identified in the rectum, transanal microsurgery may be used.

Transabdominal Resection: Abdominoperineal resection or low anterior resection or coloanal anastomosis using total mesorectal excision.

• Management Principles

- ▶ The treating surgeon should perform an endoscopy before initiating treatment
 - ▶ Removal of primary tumor with adequate margins
 - ▶ Laparoscopic surgery is not recommended outside of a clinical trial
 - ▶ Treatment of draining lymphatics by total mesorectal excision
 - ▶ Restoration of organ integrity, if possible
 - ▶ Surgery should be 5-10 weeks following full dose 5 1/2 wk neoadjuvant chemoradiation
- Total mesorectal excision
- ▶ Reduces positive radial margin rate.
 - ▶ Extend 4-5 cm below distal edge of tumors for an adequate mesorectal excision. In distal rectal cancers (ie, < 5cm from anal verge), negative distal bowel wall margin of 1-2 cm may be acceptable, this must be confirmed to be tumor free by frozen section.
 - ▶ Full rectal mobilization allows for a negative distal margin and adequate mesorectal excision.
- Lymph node dissection^{1,2}
- ▶ Biopsy or remove clinically suspicious nodes beyond the field of resection if possible.
 - ▶ Extended resection not indicated in the absence of clinically suspected nodes.

[See Criteria for Resectability of Metastases on page 2 of 3 REC-B](#)

¹Gunderson LL, Sargent DJ, Tepper JB, et al. Impact of T and N stage and treatment on survival and relapse in adjuvant rectal cancer: a pooled analysis. J Clin Oncol 2004;22(10):1785-1796.

²Greene FL, Stewart AK, Norton HJ. New tumor-node-metastasis staging strategy for node-positive (stage III) rectal cancer: an analysis. J Clin Oncol 2004;22(10):1778-1784.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURGERY (2 of 3)

CRITERIA FOR RESECTABILITY OF METASTASES AND LOCOREGIONAL THERAPIES WITHIN SURGERY

Liver

- Complete resection must be feasible based on anatomic grounds and the extent of disease, maintenance of adequate hepatic function is required.^{1,2}
- The primary tumor must have been resected for cure (R0). There should be no unresectable extrahepatic sites of disease.³⁻⁵ Plan for a debulking resection (less than an R0 resection) is not recommended.
- Patients with resectable metastatic disease and primary tumor in place should have both sites resected with curative intent. These can be resected in one operation or as a staged approach, depending on the complexity of the hepatectomy or colectomy, comorbid diseases, surgical exposure, and surgeon expertise.
- When hepatic metastatic disease is not optimally resectable based on insufficient remnant liver volume, approaches utilizing preoperative portal vein embolization or staged liver resections can be considered.
- Hepatic resection is the treatment of choice for resectable liver metastases from colorectal cancer.⁶
- Ablative techniques may be considered alone or in conjunction with resection.⁶ All original sites of disease need to be amenable to ablation or resection.
- Some institutions use intra-arterial embolization in select patients with chemotherapy resistant/refractory disease, without obvious systemic disease, with predominant hepatic metastases (category 3).
- Conformal external beam radiation therapy should not be used unless the patient is symptomatic or in the setting of a clinical trial.
- Re-resection can be considered in selected patients.⁷

Lung

- Complete resection based on the anatomic location and extent of disease with maintenance of adequate function is required.⁸⁻¹¹
- The primary tumor must have been resected for cure (R0).
- Resectable extrapulmonary metastases do not preclude resection.¹²⁻¹⁵
- Re-resection can be considered in selected patients.¹⁶
- Ablative techniques can be considered when unresectable and amenable to complete ablation.
- Patients with resectable synchronous metastases can be resected synchronously or using a staged approach.

Evaluation for conversion to resectable disease

- Re-evaluation for resection should be considered in otherwise unresectable patients after 2 months of preoperative chemotherapy and every 2 months thereafter.¹⁷⁻²⁰
- Disease with a higher likelihood of being converted to resectable are those with initially convertible disease distributed within limited sites.
- When considering whether disease has been converted to resectable, all original sites need to be amenable to resection.²¹ Preoperative chemotherapy regimens with high response rates should be considered for patients with potentially convertible disease.²²

[See footnotes on page 3 of 3 REC-B](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURGERY (3 of 3)
CRITERIA FOR RESECTABILITY OF METASTASES - REFERENCES

- ¹Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of indications for resection. Registry of Hepatic Metastases. *Surgery* 1988;103:278-288.
- ²Hughes KS, Simon R, Songhorabodi S, et al. Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of patterns of recurrence. *Surgery* 1986;100:278-284.
- ³Fong Y, Cohen AM, Fortner JG, et al. Liver resection for colorectal metastases. *J Clin Oncol* 1997;15:938-946.
- ⁴Nordlinger B, Quilichini MA, Parc R, Hannoun L, Delva E, Huguet C. Surgical resection of liver metastases from colo-rectal cancers. *Int Surg* 1987;72:70-72.
- ⁵Fong Y, Fortner J, Sun RL, Brennan MF, Blumgart LH. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: analysis of 1001 consecutive cases. *Ann Surg* 1999;230:309-318; discussion 318-321.
- ⁶Abdalla EK, Vauthey JN, Ellis LM, et al. Recurrence and outcomes following hepatic resection, radiofrequency ablation, and combined resection/ablation for colorectal liver metastases. *Ann Surg* 2004;239:818-825; discussion 825-7.
- ⁷Adam R, Bismuth H, Castaing D, et al. Repeat hepatectomy for colorectal liver metastases. *Ann Surg* 1997;225:51-62.
- ⁸McAfee MK, Allen MS, Trastek VF, Ilstrup DM, Deschamps C, Pairolero PC. Colorectal lung metastases: results of surgical excision. *Ann Thorac Surg* 1992;53:780-785; discussion 785-786.
- ⁹Regnard JF, Grunenwald D, Spaggiari L, et al. Surgical treatment of hepatic and pulmonary metastases from colorectal cancers. *Ann Thorac Surg* 1998;66:214-218; discussion 218-219.
- ¹⁰Inoue M, Kotake Y, Nakagawa K, Fujiwara K, Fukuhara K, Yasumitsu T. Surgery for pulmonary metastases from colorectal carcinoma. *Ann Thorac Surg* 2000;70:380-383.
- ¹¹Sakamoto T, Tsubota N, Iwanaga K, Yuki T, Matsuoka H, Yoshimura M. Pulmonary resection for metastases from colorectal cancer. *Chest* 2001;119:1069-1072.
- ¹²Rena O, Casadio C, Viano F, et al. Pulmonary resection for metastases from colorectal cancer: factors influencing prognosis. Twenty-year experience. *Eur J Cardiothorac Surg* 2002;21:906-912.
- ¹³Irshad K, Ahmad F, Morin JE, Mulder DS. Pulmonary metastases from colorectal cancer: 25 years of experience. *Can J Surg* 2001;44:217-221.
- ¹⁴Ambiru S, Miyazaki M, Ito H, et al. Resection of hepatic and pulmonary metastases in patients with colorectal carcinoma. *Cancer* 1998;82:274-278.
- ¹⁵Yano T, Hara N, Ichinose Y, Yokoyama H, Miura T, Ohta M. Results of pulmonary resection of metastatic colorectal cancer and its application. *J Thorac Cardiovasc Surg* 1993;106:875-879.
- ¹⁶Hendriks JM, Romijn S, Van Putte B, et al. Long-term results of surgical resection of lung metastases. *Acta Chir Belg* 2001;101:267-272.
- ¹⁷Adam R, Avisar E, Ariche A, et al. Five-year survival following hepatic resection after neoadjuvant therapy for nonresectable colorectal. *Ann Surg Oncol* 2001;8:347-353.
- ¹⁸Rivoire M, De Cian F, Meeus P, Negrier S, Sebban H, Kaemmerlen P. Combination of neoadjuvant chemotherapy with cryotherapy and surgical resection for the treatment of unresectable liver metastases from colorectal carcinoma. *Cancer* 2002;95:2283-2292.
- ¹⁹Vauthey JN, Pawlik TM, Ribero D, et al. Chemotherapy regimen predicts steatohepatitis and an increase in 90-day mortality after surgery for hepatic colorectal metastases. *J Clin Oncol*. 2006 May 1;24(13):2065-72.
- ²⁰Pawlik TM, Olin K, Gleisner AL, et al. Preoperative chemotherapy for colorectal liver metastases: impact on hepatic histology and postoperative outcome. *J Gastrointest Surg*. 2007 Jul;11(7):860-8.
- ²¹Benoist S, Brouquet A, Penna C, et al. Complete response of colorectal liver metastases after chemotherapy: does it mean cure? *J Clin Oncol*. 2006 Aug 20;24(24):3939-45.
- ²²Bartlett DL, Berlin J, Lauwers GY, et al. Chemotherapy and regional therapy of hepatic colorectal metastases: expert consensus statement. *Ann Surg Oncol*. 2006;13:1284-92.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF ADJUVANT THERAPY (1 of 2)

Adjuvant therapy for rectal cancer consists of regimens that include both concurrent chemotherapy/RT and adjuvant chemotherapy. The chemotherapy/RT may be administered either pre or postoperatively.

Postoperative adjuvant chemotherapy for patients receiving preoperative chemotherapy/RT:

- 5-FU 380 mg/m²/day on days 1-5 ± leucovorin IV 20 mg/m² on days 1-5 every 28 days x 4 cycles^{1,2}
- 5-FU 500 mg/m² IV bolus injection 1 h after the start of leucovorin infusion, once a wk for 6 wks x 3 cycles
Leucovorin 500 mg/m² IV over 2 h once a wk for 6 weeks x 3 cycles^{3,4}
 - ▶ A cycle is comprised of 6 wks followed by 2 wks of rest.

Postoperative adjuvant regimens for patients not receiving preoperative therapy:

- 5-FU + leucovorin x 1 cycle, then concurrent chemotherapy/XRT (see below for regimens), then 5-FU/leucovorin x 2 cycles^{3,4}
 - ▶ 5-FU 500 mg/m² IV bolus injection one h after the start of the leucovorin infusion, once a wk for 6 wks + leucovorin 500 mg/m² IV over 2 h once a wk for 6 wks
 - ▶ A cycle is comprised of 6 wks followed by 2 wks of rest.
- 5-FU ± leucovorin x 2 cycles, then concurrent chemotherapy/RT (see below for regimens), then 5-FU ± leucovorin x 2 cycles¹
 - ▶ 5-FU 425 mg/m²/d and leucovorin 20 mg/m²/d, days 1-5 and 29-33 before RT. After RT, the regimen is 5-FU 380 mg/m²/d and leucovorin 20 mg/m²/d for 5 consecutive days x 2 cycles
- FOLFOX (category 2B)
 - ▶ FOLFOX 4
Oxaliplatin 85 mg/m² IV over 2 hours, day 1
Leucovorin 200 mg/m² IV over 2 hours, days 1 and 2
Followed on days 1 and 2 by 5-FU 400 mg/m² IV bolus, then 600 mg/m² IV over 22 hours continuous infusion
Repeat every 2 weeks x 4 cycles⁵
 - ▶ mFOLFOX 6
Oxaliplatin 85 mg/m² IV over 2 hours, day 1
Leucovorin* 400 mg/m² IV over 2 hours, day 1
5-FU 400 mg/m² IV bolus on day 1, then 1200 mg/m²/day x 2 days (total 2400 mg/m² over 46-48 hours)** continuous infusion
Repeat every 2 weeks x 4 cycles^{6,7}
- Capecitabine⁸ (category 2B)
Capecitabine 1250 mg/m² twice daily days 1-14 every 3 wks x 24 wks

*Levo-leucovorin dose is 200 mg/m² of levo-leucovorin. The equivalent dose of leucovorin is 400 mg/m².

**NCCN recommends limiting chemotherapy orders to 24 h units (ie, 1200 mg/m²/day NOT 2400 mg/m²/day over 46 hours) to minimize medication errors.

Dosing Schedules for concurrent chemotherapy/RT:

- XRT + continuous infusion 5-FU⁹
5-FU 225 mg/m² over 24 h 7 d/wk during XRT
- XRT + 5-FU/leucovorin¹
5-FU 400 mg/m² IV bolus + leucovorin 20 mg/m² IV bolus for 4 d during wk 1 and 5 of XRT
- XRT + Capecitabine^{10,11} (category 2B)
Capecitabine 825 mg/m² twice daily 5 or 7 d/wk + XRT x 5 wks

[See footnotes on page 2 of 2 REC-C](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF ADJUVANT THERAPY (2 of 2)
REFERENCES

- ¹Tepper JE, O'Connell M, Niedzwiecki D, et al. Adjuvant therapy in rectal cancer: analysis of stage, sex, and local control--final report of Intergroup 0114. *J Clin Oncol* 2002;20:1744-1750.
- ²Sauer R, Becker H, Hohenberger W, et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med* 2004;351:1731-40.
- ³Petrelli N, Herrera L, Rustum Y et al. A prospective randomized trial of 5-fluorouracil versus 5-fluorouracil and high-dose leucovorin versus 5-fluorouracil and methotrexate in previously untreated patients with advanced colorectal carcinoma. *J Clin Oncol* 1987;5:1559-1565.
- ⁴Petrelli N, Douglass Jr HO, Herrare L, et al. The modulation of lfuorouracil with leucovorin in metastatic colorectal carcinoma: a prospective randomized phase III trial. *J Clin Oncol* 1989;7:1419-1426.
- ⁵Goldberg R, Sargent DJ, Morton RF et al. A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 2004; 22(1):23-30.
- ⁶Cheeseman S, Joel S, Chester J, et al. A "modified de Gramont" regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Brit J Cancer* 2002;87:393-399.
- ⁷Welles L, Hochster H, Ramanathan R et al. Preliminary results of a randomized study of safety and tolerability of three oxaliplatin-based regimens as first-line treatment for advanced colorectal cancer ("Tree" study). *J Clin Oncol* 2004;22(Suppl):Abstract 3537.
- ⁸Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352(26):2696-2704.
- ⁹O'Connell MJ, Martenson JA, Wieand HS, et al. Improving adjuvant therapy for rectal cancer by combining protracted-infusion fluorouracil with radiation therapy after curative surgery. *N Engl J Med* 1994; 331:502-507.
- ¹⁰Krishnan S, Janjan N, Skibber, J, et al. Phase II study of capecitabine and radiation plus concomitant boost in the treatment of locally advanced rectal cancer. *Int J Radiation Oncol Biol Phys* 2006;66:762-71.
- ¹¹Das P, Lin, E, Bhatia S, et al. Neoadjuvant Chemoradiation with Capecitabine versus Infusional 5-fluorouracil (5-FU) for Locally Advanced Rectal Cancer: a Matched Pair Analysis. *Int J Radiation Oncol Biol Phys* 2006;66:1378-83.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.


PRINCIPLES OF RADIATION THERAPY

- Radiation therapy fields should include the tumor or tumor bed, with a 2-5 cm margin, the presacral nodes, and the internal iliac nodes. The external iliac nodes should also be included for T4 tumors involving anterior structures. Consider inguinal nodes for tumors invading into the distal anal canal.
- Multiple radiation therapy fields should be used (generally a 3 or 4 field technique). Positioning and other techniques to minimize the volume of small bowel in the fields should be encouraged.
- For postoperative patients treated by abdominoperineal resection, the perineal wound should be included within the fields.
- Intensity modulated radiotherapy (IMRT) or tomotherapy should only be used in the setting of a clinical trial.
- Radiation doses:
 - ▶ 45-50 Gy in 25-28 fractions to the pelvis.
 - ▶ For resectable cancers, after 45 Gy a tumor bed boost with a 2 cm margin of 5.4 Gy in 3 fractions could be considered for preoperative radiation and 5.4-9.0 Gy in 3-5 fractions for postoperative radiation.
 - ▶ Small bowel dose should be limited to 45 Gy.
- Intraoperative radiotherapy (IORT), if available, should be considered for very close or positive margins after resection, as an additional boost, especially for patients with T4 or recurrent cancers. If IORT is not available, 10-20 Gy external beam radiation to a limited volume could be considered soon after surgery, prior to adjuvant chemotherapy.
- For unresectable cancers, doses higher than 54 Gy may be required.
- 5-fluorouracil based chemotherapy should be delivered as continuous infusion or as a bolus daily with radiation.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CONTINUUM OF CARE - CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 1 of 6)




Patient not appropriate for intensive therapy, see REC-E 2 of 6

See footnotes on page REC-E 3 of 6

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CONTINUUM OF CARE - CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 2 of 6)

Initial therapy



[See footnotes on page REC-E 3 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE (PAGE 3 of 6)

- ¹For chemotherapy references, [see Chemotherapy Regimens and References \(REC-E pages 4 - 6\)](#).
- ²Discontinuation of oxaliplatin should be strongly considered from FOLFOX or CapeOX after 3 months of therapy (or sooner if significant neurotoxicity develops > grade 3) with other drugs maintained (fluoropyrimidine + bevacizumab) until time of tumor progression. Oxaliplatin may be reintroduced if it was discontinued previously for neurotoxicity rather than disease progression. Tournigand C, Cervantes A, Figer A, et al. OPTIMOX1: A randomized study of FOLFOX4 or FOLFOX7 with oxaliplatin in a stop-and-go fashion in advanced colorectal cancer - A GERCOR Study. *J Clin Oncol* 2006;24:394-400.
- ³The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Some data suggest that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine. The relative efficacy of CapeOx with lower starting doses of capecitabine has not been addressed in large scale randomized trials. For good performance status patients, the 1000 mg/m² twice daily dose is the recommended starting dose, with close monitoring in the first cycle for toxicity, and dose adjustments as indicated.
- ⁴There are no prospective data to support continuation of bevacizumab with a second-line regimen after first progression on a bevacizumab-containing regimen and is not recommended. If bevacizumab not used in initial therapy, it may be appropriate to consider if there is no contraindication to therapy. There is an increased risk of stroke and other arterial events especially in age ≥ 65. The use of bevacizumab may interfere with wound healing.
- ⁵Combination therapy involving more than one biologic agent is not recommended. Hecht JR, Mitchell T, Chidiac C, et al. An updated analysis of safety and efficacy of oxaliplatin/bevacizumab +/- panitumumab for first-line treatment of metastatic colorectal cancer from a randomized, controlled trial (PACCE). 2008 Gastrointestinal Cancers Symposium. Abstract 273. Punt CJ, Tol J, Rodneburg J, et al randomized phase III study of capecitabine, oxaliplatin, and bevacizumab with or without cetuximab in advanced colorectal cancer, the CAIRO 2 study of the Dutch Colorectal Cancer Group. *J Clin Oncol* 28:2008 (May 20 suppl; abstract LBA4011).
- ⁶[See Principles of Pathologic Review \(REC-A\)](#) - KRAS Mutation Testing.
- ⁷Irinotecan should be used with caution and with decreased doses in patients with Gilbert's disease or elevated serum bilirubin. There is a commercially available test for UGT1A1. Guidelines for use in clinical practice have not been established.
- ⁸Infusional 5-FU is preferred. Bolus regimens of 5-FU are inappropriate as combination regimens with oxaliplatin or irinotecan.
- ⁹A treatment option for patients not able to tolerate oxaliplatin or irinotecan.
- ¹⁰Data are not mature for the addition of biologic agents to FOLFOXIRI.
- ¹¹Cetuximab is indicated in combination with irinotecan-based therapy or as single agent therapy for patients who cannot tolerate irinotecan.
- ¹²EGFR testing has no demonstrated predictive value, and therefore routine EGFR testing is not recommended. No patient should be included or excluded from cetuximab or panitumumab therapy on the basis of EGFR test results.
- ¹³There are no data, nor is there a compelling rationale, to support the use of panitumumab after clinical failure on cetuximab, or the use of cetuximab after clinical failure on panitumumab. As such, the use of one of these agents after therapeutic failure on the other is not recommended.
- ¹⁴There are no data to support the combination of panitumumab with chemotherapy.
- ¹⁵Single agent or combination therapy with capecitabine, mitomycin, or gemcitabine has not been shown to be effective in this setting.
- ¹⁶Patients with diminished creatinine clearance may require dose modification of capecitabine.
- ¹⁷Routine use of bevacizumab + cetuximab is not recommended in patients with prior bevacizumab progression.
- ¹⁸The use of single agent capecitabine as a salvage therapy after failure on a fluoropyrimidine-containing regimen has been shown to be ineffective, and this is therefore not recommended.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE (PAGE 4 of 6)

FOLFOX**FOLFOX 4**

Oxaliplatin 85 mg/m² IV over 2 hours, day 1
Leucovorin 200 mg/m² IV over 2 hours, days 1 and 2
Followed on days 1 and 2 by 5-FU 400 mg/m² IV bolus, then 600 mg/m² IV over 22 hours continuous infusion
Repeat every 2 weeks¹

mFOLFOX 6

Oxaliplatin 85 mg/m² IV over 2 hours, day 1
Leucovorin* 400 mg/m² IV over 2 hours, day 1
5-FU 400 mg/m² IV bolus on day 1, then 1200 mg/m²/day x 2 days (total 2400 mg/m² over 46-48 hours)[†] continuous infusion
Repeat every 2 weeks^{2,3}

CapeOX^{3,4}

Oxaliplatin 130 mg/m² day 1, Capecitabine 850-1000[‡] mg/m² twice daily for 14 days
Repeat every 3 weeks

FOLFIRI^{5,6}

Irinotecan 180 mg/m² IV over 30-120 minutes, day 1
Leucovorin 200 mg/m² IV infusion to match duration of irinotecan infusion, days 1 and 2
Followed on days 1 and 2 by 5-FU 400 mg/m² IV bolus, then 600 mg/m² IV over 22 hours continuous infusion
Repeat every 2 weeks

Irinotecan 180 mg/m² IV over 30-120 minutes, day 1
Leucovorin 400* mg/m² IV infusion to match duration of irinotecan infusion, day 1
5-FU 400 mg/m² IV bolus day 1, then 1200 mg/m²/day x 2 days (total 2400 mg/m² over 46-48 hours)[†] continuous infusion
Repeat every 2 weeks

Bevacizumab + 5-FU containing regimens:^{7,8,9}

Bevacizumab 5 mg/kg IV every 2 weeks +
5-FU and Leucovorin
or FOLFOX¹⁰
or FOLFIRI
Bevacizumab 7.5 mg/kg IV every 3 weeks + CapeOX⁴

*Levoleucovorin dose is 200 mg/m². The equivalent dose of leucovorin is 400 mg/m².

[†]NCCN recommends limiting chemotherapy orders to 24 h units (ie, 1200 mg/m²/day NOT 2400 mg/m²/day over 46 hours) to minimize medication errors.

[‡]The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine. The relative efficacy of CapeOx with lower starting doses of capecitabine has not been addressed in large scale randomized trials.

[See footnotes on page 6 of 6 REC-E](#)

[See Additional Chemotherapy Regimens 5 of 6 REC-E](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE (PAGE 5 of 6)

CHEMOTHERAPY REGIMENS

Capecitabine¹¹

2000-2500 mg/m²/day PO in two divided doses, days 1-14,
followed by 7 days rest
Repeat every 3 weeks

**Bolus or infusional 5-FU/leucovorin
Roswell-Park regimen¹²**

Leucovorin 500 mg/m² IV over 2 hours, days 1, 8, 15, 22, 29, and 36
5-FU 500 mg/m² IV bolus 1 hour after start of Leucovorin,
days 1, 8, 15, 22, 29, 36
Repeat every 8 weeks

Biweekly¹³

Leucovorin 200 mg/m² IV over 2 hours, days 1 and 2
5-FU 400 mg/m² IV bolus, then 600 mg/m² IV over 22 hours
continuous infusion, days 1 and 2
Repeat every 2 weeks

Simplified biweekly infusional 5-FU/LV (sLV5FU2)¹⁴

Leucovorin 400* mg/m² IV over 2 hours on day 1,
followed by 5-FU bolus 400 mg/m² and then 1200 mg/m²/day x 2
days (total 2400 mg/m² over 46-48 hours)[†] continuous infusion
Repeat every 2 weeks

Weekly

Leucovorin 20 mg/m² as a 2 h infusion
5-FU 500 mg/m² bolus administered 1 h after LV infusion
Repeat every week¹⁵
5-FU 2600 mg/m² by 24 h infusion plus leucovorin 500 mg/m²
Repeat every week¹⁶

*Levoleucovorin dose is 200 mg/m². The equivalent dose of leucovorin is 400 mg/m².

[†]NCCN recommends limiting chemotherapy orders to 24 h units (ie, 1200 mg/m²/day NOT 2400 mg/m²/day over 46 hours) to minimize medication errors.

FOLFOXIRI¹⁷

Irinotecan 165 mg/m² IV day 1, oxaliplatin 85 mg/m² day 1,
leucovorin 200 mg/m² day 1, fluorouracil 3,200 mg/m² 48
continuous infusion starting on day 1
Repeat every 2 weeks

Irinotecan^{18,19}

Irinotecan 125 mg/m² IV over 30-90 minutes, days 1, 8, 15, 22
Repeat every 6 weeks

Irinotecan 300-350 mg/m² IV over 30-90 minutes, day 1
Repeat every 3 weeks

Cetuximab (KRAS wild-type gene only) ± irinotecan²⁰
Cetuximab 400 mg/m² 1st infusion, then 250 mg/m² IV weekly
or
Cetuximab 500 mg/m² IV every 2 weeks²¹

±
Irinotecan 300-350 mg/m² IV every 3 weeks
or
Irinotecan 180 mg/m² IV every 2 weeks
or
Irinotecan 125 mg/m² every week for 4 weeks
Every 6 weeks

Cetuximab (KRAS wild-type gene only)
Cetuximab 400 mg/m² 1st infusion, then 250 mg/m² IV weekly

Panitumumab²² (KRAS wild-type gene only)
Panitumumab 6 mg/kg IV over 60 minutes every 2 weeks

[See footnotes on page 6 of 6 REC-E](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

CHEMOTHERAPY FOR ADVANCED OR METASTATIC DISEASE (PAGE 6 of 6)

CHEMOTHERAPY REFERENCES

- ¹Goldberg R, Sargent DJ, Morton RF et al. A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 2004;22:23-30.
- ²Cheeseman S, Joel S, Chester J, et al. A “modified de Gramont” regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Brit J Cancer* 2002;87:393-399.
- ³Cassidy J, Clarke S, Diaz Rubio E, et al. First efficacy and safety results from Xelox-1/NO16966, a randomized 2 x 2 factorial phase III trial of Xelox vs Folfox4 + bevacizumab or placebo in first-line metastatic colorectal cancer. *Ann Oncol*;17(suppl 9):late breaking abstract #3.
- ⁴European studies showing equivalent efficacy for CapeOX used at a higher dose; however, European patients consistently tolerate capecitabine with less toxicity than American patients.
- ⁵Douillard J, Cunningham D, Roth A et al. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *The Lancet* 2000;355:1041-1047.
- ⁶Andre T, Louvet C, Maindault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
- ⁷Kabbinavar FF, Hambleton J, Mass RD, et al. Combined analysis of efficacy: the addition of bevacizumab to fluorouracil/leucovorin improves survival for patients with metastatic colorectal cancer. *J Clin Oncol*. 2005;23:3706-3712.
- ⁸Hurwitz HI, Fehrenbacher L, Hainsworth JD, et al. Bevacizumab in combination with fluorouracil and leucovorin: an active regimen for first-line metastatic colorectal cancer. *J Clin Oncol*. 2005;23:3502-3508.
- ⁹Reidy DL, Chung KY, Timoney JP, et al. Bevacizumab 5 mg/kg can be infused safely over 10 minutes. *J Clin Oncol* 2007;25:2691-2695.
- ¹⁰Giantonio BJ, Catalano PJ, Meropol NJ, et al. High-dose bevacizumab in combination with FOLFOX4 improves survival in patients with previously treated advanced colorectal cancer: Results from the Eastern Cooperative Oncology Group (ECOG) study E3200. 2005 ASCO Gastrointestinal Cancers Symposium;Abstract 169a.
- ¹¹VanCutsem E, Twelves C, Cassidy J, et al. Oral capecitabine compared with intravenous fluorouracil plus leucovorin in patients with metastatic colorectal cancer: results of a large phase III study. *J Clin Oncol* 2001;19:4097-4106.
- ¹²Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from National Surgical Adjuvant Breast and Bowel Protocol C-03. *J Clin Oncol* 1993;11:1879-1887.
- ¹³de Gramont A, Bosset JF, Milan C, et al. Randomized trial comparing monthly low-dose leucovorin and fluorouracil bolus with bimonthly high-dose leucovorin and fluorouracil bolus plus continuous infusion for advanced colorectal cancer: a French intergroup study. *J Clin Oncol* 1997;15:808-815.
- ¹⁴Andre T, Louvet C, Maindault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-FU fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35:1343-7.
- ¹⁵Jäger E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. *J Clin Oncol* 1996;14:2274-2279.
- ¹⁶Douillard JY, Cunningham D, Roth AD, et al. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *The Lancet* 2000;355:1041-47.
- ¹⁷Falcone A, Ricci S, Brunetti I, et al. Phase III trial of infusional fluorouracil, leucovorin, oxaliplatin, and irinotecan (FOLFOXIRI) compared with infusional fluorouracil, leucovorin, and irinotecan (FOLFIRI) as first-line treatment for metastatic colorectal cancer: The Gruppo Oncologico Nord Ovest. *J Clin Oncol* 2007;25(13):1670-1676.
- ¹⁸Cunningham D, Pyrhonen S, James R, et al. Randomised trial of irinotecan plus supportive care versus supportive care alone after fluorouracil failure for patients with metastatic colorectal cancer. *The Lancet* 1998;352:1413-1418.
- ¹⁹Fuchs CS, Moore MR, Harker G, et al. Phase III comparison of two irinotecan dosing regimens in second-line therapy of metastatic colorectal cancer. *J Clin Oncol* 2003;21:807-814.
- ²⁰Cunningham D, Humblet Y, Siena S, et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* 2004;351:337-345.
- ²¹Van Cutsem E, Humblet H, Gelderblom J, et al. Cetuximab dose-escalation in patients with metastatic colorectal cancer with no or slight skin reactions on cetuximab standard dose treatment (EVEREST): Pharmacokinetic and efficacy data of a randomized study. 2007 Gastrointestinal Cancers Symposium. Abstract 237.
- ²²Van Cutsem E, Peeters M, Siena S, et al. Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol* 2007;25:1658-1664.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURVIVORSHIP
Colorectal Long-term Follow-up Care (1 of 3)**CRC Cancer Surveillance:**

- History and Physical every 3-6 months for 2 years, then every 6 months for 3 years.
- CEA every 3-6 months for 2 years, then every 6 months for 3 years.
- CT scan of abdomen and pelvis annually for 3 years.
- Colonoscopy at 1 year, then as clinically indicated.

Cancer Screening Recommendations:• **Breast Cancer:**

- ▶ Periodic self breast exam (SBE) encouraged (optional)
- ▶ Clinical breast exam (CBE) every 1-3 years between ages 20 and 40
- ▶ Annual mammogram with clinical breast exam beginning at age 40.
- ▶ Women at high risk (greater than 20% lifetime risk) should get breast MRI and mammogram annually.
- ▶ See [NCCN Breast Cancer Screening and Diagnosis Guidelines](#)

• **Cervical Cancer:**

- ▶ Annual cervical cytology testing with conventional smears or every 2 years with liquid-based cytology for women up to age 30.
- ▶ After age 30, screening may be every 2-3 years if 3 negative/satisfactory annually cervical cytology tests documented.
- ▶ Alternatively, human papilloma virus (HPV) DNA testing for women age 30 and over, combined with cervical cytology.
- ▶ If cervical cytology and HPV DNA testing both negative, testing may be performed every 3 years.
- ▶ Counseling regarding HPV infection.
- ▶ Women over age 70 with no abnormal testing in last 10 years and 3 normal tests in a row may discontinue screening.
- ▶ Women without a cervix from a total abdominal hysterectomy do not need to be screened.
- ▶ See [NCCN Cervical Cancer Screening Guidelines](#)

• **Prostate Cancer:**

- ▶ Annual prostate specific antigen (PSA) testing and digital rectal exam (DRE) beginning at age 50
- ▶ For high risk men (African-American males and those with a family history of prostate cancer): PSA testing and DRE beginning at age 40.
- ▶ See [NCCN Prostate Cancer Early Detection Guidelines](#)

[Continued](#)

¹American Cancer Society Guidelines for Early Detection of Cancer:

http://www.cancer.org/docroot/PED/content/PED_2_3X_ACS_Cancer_Detection_Guidelines_36.asp, Accessed September 21, 2008.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURVIVORSHIP
Colorectal Long-term Follow-up Care (2 of 3)**Management of Late Sequelae of Disease or Treatment:²⁻⁶**

- **Chronic Diarrhea or Incontinence**
 - ▶ Consider anti-diarrheal agents, bulk-forming agents, diet manipulation, and protective undergarments.
- **Oxaliplatin-Induced Neuropathy**
 - ▶ Consider the use of gabapentin and/or tricyclic antidepressants for persistent, painful neuropathy.
- **Bone Health After Pelvic Radiation**
 - ▶ Consider monitoring of bone density or evaluation for pelvic fractures with pelvic pain if previously received pelvic radiation
- **Sexual Dysfunction After Pelvic Radiation**
 - ▶ Screen for erectile dysfunction and dyspareunia in those who received pelvic radiation
 - ▶ Consider referral to urologist or gynecologist for persistent symptoms.

Immunizations:⁷

- Annual trivalent inactivated influenza vaccination
- Pneumococcal vaccination with revaccination as appropriate

Routine Health Monitoring and Screening:

- Cholesterol, blood pressure, and glucose monitoring
- Bone density testing as appropriate
- Routine dental examinations
- Routine sun protection
- Screening for depression as appropriate

[Continued](#)

²Schneider EC, Malin JL, Kahn KL, et al. Surviving colorectal cancer. *Cancer* 2007;110: 2075-2082.

³Sprangers MAG, Taal BG, Aaronson NK, et al. Quality of life in colorectal cancer: stoma vs. nonstoma patients. *Dis Colon Rectum* 1995;38:361-369.

⁴Kalso E, Tasmuth T, Neuvonen PJ. Amitriptyline effectively relieves neuropathic pain following treatment of breast cancer. *Pain* 1995;64: 293-302.

⁵Caraceni A, Zecca E, Bonezzi C, et al. Gabapentin for neuropathic cancer pain: a randomized controlled trial from the Gabapentin Cancer Pain Study Group. *J Clin Oncol* 2004;22: 2909-2917.

⁶Baxter NN, Habermann EB, Tepper JE, et al. Risk of pelvic fractures in older women following pelvic irradiation. *JAMA* 2005; 294: 2587-2593.

⁷Advisory Committee on Immunization Practices. Recommended adult immunization schedule: United States, October 2007–September 2008. *Ann Intern Med*. 2007;147:725-9.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURVIVORSHIP
Colorectal Long-term Follow-up Care (3 of 3)**Counseling Regarding Healthy Lifestyle and Wellness:**⁸⁻¹¹

- Screening and counseling to maintain a healthy weight.
- Screening for physical activity and counseling to adopt a physically active lifestyle (Recommended activity: at least 30 minutes or more of moderate to vigorous physical activity at least 5 days of the week).
- Screening and counseling for alcohol use.
- Screening and counseling for tobacco use with emphasis on smoking cessation.
- Counseling regarding healthy diet adoption, with emphasis on plant sources.

Prescription for Survivorship and Transfer of Care to Primary Care Physician:¹²

- Include overall summary of treatment, including all surgeries, radiation treatments, and chemotherapy received
- Describe possible clinical course, including expected time to resolution of acute toxicities, long-term effects of treatment, and possible late sequelae of treatment
- Include surveillance recommendations
- Delineate appropriate timing of transfer of care with specific responsibilities identified for PCP and Oncologist.

⁸American Cancer Society Guidelines on Nutrition and Physical Activity for Cancer Prevention, http://www.cancer.org/docroot/PED/content/PED_3_2X_Diet_and_Activity_Factors_That_Affect_Risks.asp?sitearea=PED, Accessed September 21, 2008.

⁹Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. J Clin Oncol 2006;24:3535-3541.

¹⁰Meyerhardt JA, Niedzwiecki D, Hollis D, et al. Association of dietary patterns with cancer recurrence and survival in patients with stage III colon cancer. JAMA 2007;298:754-764.

¹¹Dignam JL, Polite BN, Yothers G, et al. Body Mass Index and outcomes in patients who receive adjuvant chemotherapy for colon cancer. J Natl Cancer Inst 2006;98:1647-54.

¹²Hewitt M, Greenfield S, Stovall E. From Cancer Patient to Cancer Survivor: Lost in Transition. Washington, D.C.:The National Academies Press;2006.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

Staging

Table 1

American Joint Committee on Cancer (AJCC) TNM Staging System for Colorectal Cancer* Rectal Cancer < 12 cm

Primary Tumor (T)

- TX Primary tumor cannot be assessed
- T0 No evidence of primary tumor
- Tis Carcinoma in situ: intraepithelial or invasion of lamina propria†
- T1 Tumor invades submucosa (SM 1-3)
- T2 Tumor invades muscularis propria
- T3 Tumor invades through the muscularis propria into the subserosa, or into nonperitonealized pericolic or perirectal tissues
- T4 Tumor directly invades other organs or structures, and/or perforates visceral peritoneum‡

Regional Lymph Nodes (N)§

- NX Regional lymph nodes cannot be assessed
- N0 No regional lymph node metastasis
- N1 Metastasis in 1 to 3 regional lymph nodes
- N2 Metastasis in 4 or more regional lymph nodes

Distant Metastasis (M)

- MX Distant metastasis cannot be assessed
- M0 No distant metastasis
- M1 Distant metastasis

Stage Grouping

Stage	T	N	M	Dukes¶	MAC¶
0	Tis	N0	M0	-	-
I	T1	N0	M0	A	A
	T2	N0	M0	A	B1
IIA	T3	N0	M0	B	B2
IIB	T4	N0	M0	B	B3
IIIA	T1-T2	N1	M0	C	C1
IIIB	T3-T4	N1	M0	C	C2/C3
IIIC	Any T	N2	M0	C	C1/C2/C3
IV	Any T	Any N	M1	-	D

Histologic Grade (G)

- GX Grade cannot be assessed
- G1 Well differentiated
- G2 Moderately differentiated
- G3 Poorly differentiated
- G4 Undifferentiated

*Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the *AJCC Cancer Staging Manual, Sixth Edition (2002)* published by Springer-Verlag New York. (For more information, visit www.cancerstaging.net.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed written permission of Springer-Verlag New York on behalf of the AJCC.

†Tis includes cancer cells confined within the glandular basement membrane (intraepithelial) or lamina propria (intramucosal) with no extension through the muscularis mucosae into the submucosa.

‡Direct invasion in T4 includes invasion of other segments of the colorectum by way of the serosa; for example, invasion of the sigmoid colon by a carcinoma of the cecum. Tumor that is adherent to other organs or structures, macroscopically, is classified T4. However, if no tumor is present in the adhesion, microscopically, the classification should be pT3. The V and L substaging should be used to identify the presence or absence of vascular or lymphatic invasion.

§A tumor nodule in the pericorectal adipose tissue of a primary carcinoma without histologic evidence of residual lymph node in the nodule is classified in the pN category as a regional lymph node metastasis if the nodule has the form and smooth contour of a lymph node. If the nodule has an irregular contour, it should be classified in the T category and also coded as V1 (microscopic venous invasion) or as V2 (if it was grossly evident), because there is a strong likelihood that it represents venous invasion.

¶Dukes B is a composite of better (T3 N0 M0) and worse (T4 N0 M0) prognostic groups, as is Dukes C (Any TN1 M0 and Any T N2 M0). MAC is the modified Astler-Coller classification.

Note: The y prefix is to be used for those cancers that are classified after pretreatment, whereas the r prefix is to be used for those cancers that have recurred.

Discussion

This discussion is being updated to correspond with the newly updated algorithm. Last updated 10/28/08

NCCN Categories of Evidence and Consensus

Category 1: The recommendation is based on high-level evidence (e.g. randomized controlled trials) and there is uniform NCCN consensus.

Category 2A: The recommendation is based on lower-level evidence and there is uniform NCCN consensus.

Category 2B: The recommendation is based on lower-level evidence and there is nonuniform NCCN consensus (but no major disagreement).

Category 3: The recommendation is based on any level of evidence but reflects major disagreement.

All recommendations are category 2A unless otherwise noted.

Overview

In 2008 an estimated 40,740 new cases of rectal cancer will occur in the United States (23,490 cases in men; 17,250 cases in women). During the same year, it is estimated that 49,960 people will die from rectal and colon cancer.¹ Although colorectal cancer is ranked as the third most frequently diagnosed cancer in men and women in the U.S., mortality from colorectal cancer has decreased during the past 30 years. This decrease may be due to both earlier diagnosis through screening and better treatment modalities.

The recommendations in these clinical practice guidelines are classified as category 2A except where noted, meaning that there is uniform NCCN consensus, based on lower-level evidence (including clinical experience), that the recommendation is appropriate. The panel unanimously endorses patient participation in a clinical trial over

standard or accepted therapy. This is especially true for cases of advanced disease and for patients with locally aggressive colorectal cancer who are receiving combined modality treatment. The clinical practice guidelines for managing rectal cancer overlap considerably with the [NCCN Colon Cancer Guidelines](#). First-degree relatives of patients with newly diagnosed adenomas² or invasive carcinoma³ are at increased risk for colorectal cancer. Therefore, rectal cancer patients, especially those 50 years or younger, should be counseled regarding their family history as outlined in the [NCCN Colorectal Screening Guidelines](#).

TNM Staging

The NCCN Rectal Cancer Guidelines adhere to the current TNM staging system as included in the 6th edition of the American Joint Committee on Cancer's (AJCC) *Cancer Staging Manual* ([Table 1](#)).^{4,5} Stage I rectal cancer is defined as T1-T2, N0, M0. Stage II disease is subdivided into IIA (if the primary tumor is T3, N0, M0) and IIB (for T4, N0, M0 lesions). Stage III disease is subdivided into IIIA (T1-2, N1, M0), IIIB (T3-4, N1, M0), and IIIC (any T, N2, M0). Stage IV disease is defined as any T, any N, and the presence of one or more distant metastases (M1). The difference between N1 and N2 disease is the number of nodes involved: N1 lesions have 1 to 3 positive regional lymph nodes, whereas N2 tumors have 4 or more regional lymph nodes. In this version of the staging system, smooth metastatic nodules in the pericolic or perirectal fat are considered lymph node metastases and should be included in N staging. Irregularly contoured metastatic nodules in the peritumoral fat are considered vascular invasion. In addition, the 6th edition of the AJCC staging manual⁶ includes the suggestion that the surgeon mark the area of the specimen with the deepest tumor penetration so that the pathologist can directly evaluate the status of the resection margins. The surgeon is encouraged to score the completeness of the resection as (1) R0 for complete tumor resection with all margins negative; (2) R1 for incomplete tumor

resection with microscopic involvement of a margin; and (3) R2 for incomplete tumor resection with gross residual tumor that was not resected.

Pathology

Pathologic staging information is provided by examination of the surgical specimen. Some of the information that should be detailed in the report of the pathologic evaluation of rectal cancer includes: 1) gross description of the tumor and specimen 2) grade of the cancer; 3) depth of penetration and extension to adjacent structures (T); 4) number of regional lymph nodes evaluated and 5) number of positive regional lymph nodes (N); 6) the presence of distant metastases to other organs, the peritoneum of an abdominal structure, or non-regional lymph nodes (M) and 7) the status of proximal, distal, and circumferential (radial) margins^{5,7}. The prefixes “p” and “yp” used in TNM staging denote pathologic staging and pathologic staging following neoadjuvant therapy, respectively.⁸

The circumferential margin or circumferential resection margin (CRM) is an important pathologic staging parameter in rectal cancer. Whereas the radial margin for resected segments of the colon that are completely encased by a peritonealized (serosal) surface is also referred to as the peritoneal margin, the CRM is very important in segments of the colon or rectum that are either not encased or only partially encased in peritoneum.⁵ The CRM is the closest radial margin between the deepest penetration of the tumor and the edge of resected soft tissue around the rectum (ie, the retroperitoneal or subperitoneal aspect of the tumor) and should be measured in millimeters. Identification of the CRM is determined through evaluation of the outer circumference of the rectal and mesorectal specimen which often requires inking of the outer surfaces and “bread-loaf” slicing of the specimen.⁹ A positive CRM has been defined as tumor within 1-2 mm from the transected margin.¹⁰⁻¹³ Accurate pathologic assessment of the

CRM of resected rectal tumor specimens is very important since the CRM has been shown to be a strong predictor of both local recurrence and overall survival, and is an important consideration when post-operative treatment decisions are made.^{8,14,15} Furthermore, in a retrospective study of over 17,000 patients with rectal cancer, CRM was found to be a better predictor of local recurrence for patients who had received preoperative therapy when these patients were compared with patients undergoing surgery as initial therapy.¹⁶

The AJCC and College of American Pathologists (CAP) recommend evaluation of a minimum of 12 lymph nodes to accurately identify stage II colorectal cancers.^{5,6} The number of lymph nodes retrieved can vary with age of the patient, gender, and tumor grade or site.^{17,18} The extent and quality of surgical resection and pathologic review of the specimen can also have an impact on the node harvest.¹⁹ The literature lacks consensus regarding the minimal number of lymph nodes needed to accurately identify stage II rectal cancer. Most of these studies have combined rectal and colon cancers and reflect those cases with surgery as the initial treatment. Two studies confined only to rectal cancer have reported 14 and >10 lymph nodes as the minimal number to accurately identify stage II rectal cancer.^{20,21} Furthermore, the mean number of lymph nodes retrieved from rectal cancers treated with neoadjuvant therapy is significantly less than those treated by surgery alone (13 vs 19, $P < 0.05$; 7 vs 10, $P \leq 0.0001$).^{22,23} A recent retrospective analysis of data from patients with T3/T4 and/or lymph node-positive rectal cancer enrolled in the Intergroup 0114 trial showed lymph node ratio (LNR), the number of positive lymph nodes divided by the total number, to be a strong predictor of survival.²⁴ Nevertheless, the panel does not consider determination of LNR to be a substitute for an adequate lymph node evaluation.

Results of studies evaluating the sentinel node for micrometastatic disease through use of hematoxylin and eosin (H&E) staining to identify

small foci of tumor cells, or identification of particular tumor antigens through immunohistochemical (IHC) analysis have been reported.^{25,26} Although results of some of these studies seem promising, there is no uniformity in the definition of “true” clinically relevant metastatic carcinoma. Some studies have considered detection of single cells by IHC as well as isolated tumor cells (ITC) to be micrometastasis.^{27,28} In addition, results of one study demonstrated that, following neoadjuvant radiotherapy for rectal cancer, the sensitivity for the sentinel node procedure was only 40%.²⁹ Presently, the use of sentinel lymph nodes and detection of cancer cells by IHC alone should be considered investigational and the results should be used with caution in clinical management decisions.

Clinical Presentation and Treatment

Management of Polypoid Cancer

Before making a decision about surgical resection for an endoscopically resected adenomatous polyp or villous adenoma, physicians should review pathology and consult with the patient.³⁰ A malignant rectal polyp is defined as one with cancer invading through the muscularis mucosae and into the submucosa (pT1). Conversely, polyps classified as carcinoma in situ (pTis) have not penetrated into the submucosa and are therefore not considered to be capable of regional nodal metastasis.⁵ The panel recommends marking the cancerous polyp site at the time of colonoscopy or within 2 weeks. In patients with invasive cancer and adenoma (tubular, tubulovillous or villous), no additional surgery is required for pedunculated or sessile polyps, if the polyp has been completely resected with favorable histological features.³⁰ Favorable histological features include lesions of grade 1 or 2, no angiolymphatic invasion and a negative resection margin. However, in addition to the option of observation, the panel includes the option of colectomy in patients with a completely-removed, single-specimen, sessile polyp with favorable histological features and clear margins because it has been reported that patients with sessile polyps have a

10% risk of lymph node metastases.³¹ For pedunculated and sessile polyps, unfavorable histopathological features are: grade 3 or 4, angiolymphatic invasion, or a positive margin of resection. It should be noted that there is currently no consensus as to the definition of what constitutes a positive margin of resection. A positive margin has been defined as the presence of tumor within 1-2 mm from the transected margin and the presence of tumor cells within the diathermy of the transected margin.^{30,32-34} For a pedunculated or sessile polyp with fragmented specimen or margins that cannot be assessed or unfavorable pathology, either a transanal excision or a transabdominal resection is recommended (See section on [Surgical Approaches](#) used in the management of rectal cancer appropriate for resection). Results from a preoperative endoscopic ultrasound evaluation may provide additional information to guide choice of surgical approach, although the accuracy of this method to detect residual cancer is limited (see section on [Clinical Evaluation/Staging](#)).³⁵ All patients who have resected polyps should undergo total colonoscopy to rule out other synchronous polyps, as well as appropriate follow-up surveillance endoscopy.³⁶

Management of Rectal Cancer

Rectal cancer has been defined as a cancerous lesion located within 12 cm of the anal verge by rigid proctoscopy.³⁷ Some support for this definition comes from the study of Kapiteijn et al.³⁸ which included a subgroup analysis of the risk of recurrence of rectal cancer based on tumor location. Univariate analyses indicated that local recurrence rates were low for patients who had tumors with an inferior margin of 10.1 cm or more from the anal verge, and that no significant differences between patients in this group receiving radiotherapy and surgery were observed when they were compared to those undergoing surgery alone.

Determination of an optimal treatment plan for an individual patient with rectal cancer is a complex process. In addition to decisions relating to the intent of rectal cancer surgery (ie, curative or palliative), consideration must also be given to the likely functional results of treatment, including the probability of maintaining or restoring normal bowel function/anal continence, and preserving genitourinary functions. For patients with distal rectal cancer, in particular, the simultaneous achievement of the goals of cure and minimal impact on quality of life can be challenging.³⁹ Furthermore, the risk of pelvic recurrence is higher in patients with rectal cancer compared to those with colon cancer, and locally recurrent rectal cancer has frequently been associated with a poor prognosis.^{40,41} Careful patient selection with respect to particular treatment options and the use of sequenced multimodality therapy for selected patients which combines chemoradiation (chemoRT) with operative treatment as part of the treatment regimen is recommended.

Clinical Evaluation/Staging

The initial clinical workup of patients with rectal cancer provides important preoperative information on the clinical stage of disease. Since the clinical stage of the disease is used to direct decisions regarding choice of primary treatment, including surgical intent (eg, curative or palliative) and approaches, and whether to recommend preoperative chemoRT, the implications of either clinically understaging or over-staging rectal cancer can be substantial.

Patients who present with rectal cancer appropriate for resection require complete staging evaluation, including total colonoscopy to evaluate for synchronous lesions or other pathologic conditions of the colon and rectum, proctoscopy to provide a determination of the location of the cancer (eg, measurement of the distance of the tumor from the anal verge should be performed by the responsible surgeon using rigid proctoscopy), and a complete physical examination,

including assessment of performance status, to determine operative risk, carcinoembryonic antigen (CEA) determination, and baseline computed tomographic (CT) scans of the chest, abdomen and pelvis. The consensus of the panel is that a positron emission tomography (PET) scan is not routinely indicated at baseline in the absence of evidence of synchronous metastatic disease. In addition, the accessibility of rectal cancer to evaluation by certain imaging modalities, such as endoscopic ultrasound and magnetic resonance imaging (MRI), makes possible preoperative assessments of depth of tumor penetration and the presence of local lymph nodal metastases.⁴² Additional information regarding the extent of disease and the occurrence of distant metastases can be determined preoperatively through CT scans. Thus, endorectal ultrasound or endorectal or pelvic MRI, and CT scans of the chest, abdomen and pelvis are recommended for the preoperative staging of rectal cancer.

Results from a meta-analysis of 90 studies involving the accuracy of endoscopic ultrasound, MRI, and CT in preoperatively staging rectal cancer demonstrated that endoscopic ultrasound and MRI have similarly high sensitivities for evaluating the depth of tumor penetration into the muscularis propria (94%), although endoscopic ultrasound was found to be more specific than MRI in the evaluation of local tumor invasion (86% vs. 69%).⁴³ Only a very limited number of studies using CT for the purpose of T-staging have been performed, and it is not currently considered to be an optimal method for staging the extent of tumor penetration.^{43,44} Accurate assessment of nodal status is one of the greatest challenges in the preoperative staging of rectal cancer. In the meta-analysis of Bipat et al.,⁴³ the sensitivities and specificities of the 3 imaging modalities for accurately evaluating lymph node involvement were: CT (55% and 74%); endoscopic ultrasound (67% and 78%); and MRI (66% and 76%). Results from another recent meta-analysis of 84 articles, indicated that none of the 3 imaging modalities were significantly superior to another method with respect to an

accurate determination of tumor N-stage.⁴⁵ Disadvantages of endoscopic ultrasound and MRI include a high degree of operator dependence.⁴³ An advantage of MRI is its ability to provide accurate images of soft tissue structures in the mesorectum, including the mesorectal fascia. Hence, MRI evaluation of patients with more advanced rectal cancer has the potential to provide information useful in the prediction of the CRM prior to radical surgery.⁴⁴⁻⁴⁶

Clinical staging is also based on histopathologic examination of the specimen obtained via biopsy or local excision (eg, excised polyps). Endoscopic biopsy specimens of the lesion should undergo careful pathology review for evidence of invasion into the muscularis mucosa. If removal of the rectum is contemplated, early consultation with an enterostomal therapist is recommended for preoperative marking of the site and patient teaching purposes.

Surgical Approaches

A variety of surgical approaches, depending on the location and extent of disease, are used to treat the primary rectal cancer lesion.⁴⁷ These methods include local procedures, such as polypectomy, transanal excision and transanal microsurgery, and radical procedures involving an transabdominal resection (eg, low anterior resection [LAR], total mesorectal excision [TME] with coloanal anastomosis or abdominoperineal resection [APR]).

Transanal excision may be appropriate for selected early-stage cancers. Small (<3 cm), well to moderately differentiated T1 or T2 tumors (category 2B for T2 tumors) that are within 8 cm of the anal verge and limited to less than 30% of the rectal circumference, and for which there is no evidence of nodal involvement (category 2A) can be approached with transanal excision with negative margins. Transanal endoscopic microsurgery (TEM) can facilitate excision of small tumors through the anus that are located higher up in the rectum. Both transanal excision and TEM involve a full thickness excision performed

perpendicularly through the bowel wall into the perirectal fat. Negative (> 3 mm) deep and mucosal margins are required. Tumor fragmentation should be avoided. The excised specimen should be oriented and pinned before fixation and brought to the pathologist by the surgeon (ie, to facilitate an oriented histopathologic evaluation of the specimen). Advantages of a local procedure include minimal morbidity (eg, a sphincter-sparing procedure) and mortality and rapid postoperative recovery.^{39,48} If pathologic examination reveals adverse features such as high grade, positive margins, lymphovascular invasion (LVI) or perineural invasion, a more radical resection is recommended. Caution is recommended when local excision is considered in the treatment of patients with T2 tumors since data on long-term patient outcomes, including risk of local recurrence, are limited.⁴⁸

Patients with rectal cancer who do not meet requirements for local surgery should be treated with a transabdominal resection. Organ-preserving procedures which maintain sphincter function are preferable, but not possible, in all cases. For lesions in the mid to upper rectum, a low anterior resection (LAR) extended 4-5 cm below distal edge of tumor, followed by creation of a colorectal anastomosis, is the treatment of choice. Where creation of an anastomosis is not possible, colostomy is required.

Data from randomized studies evaluating use of laparoscopic surgery in the treatment of patients with rectal cancer are limited.^{49,50} In the CLASICC trial comparing laparoscopically-assisted resection to open resection, nearly half of the 794 patients were diagnosed with rectal cancer.⁴⁹ No significant differences in local recurrence, DFS, or overall survival were observed between the 2 groups of patients with rectal cancer based on surgical approach. However, factors which may confound conclusions drawn from randomized studies comparing open surgery to laparoscopically-assisted surgery for colorectal cancer have

been described,⁵¹ and laparoscopic surgery for rectal cancer is not recommended by the panel outside of a clinical trial.

For low rectal lesions, abdominoperineal resection (APR) or total mesorectal excision (TME) with coloanal anastomosis is required. A TME involves an en bloc removal of the mesorectum, including associated vascular and lymphatic structures, fatty tissue, and mesorectal fascia as a “tumor package” through sharp dissection and is designed to spare the autonomic nerves.^{39,52} In cases where anal function is intact and distal clearance is adequate, the TME may be followed by creation of a coloanal anastomosis. An APR involves en bloc resection of the rectosigmoid, the rectum, and the anus, as well as the surrounding mesentery, mesorectum, and perianal soft tissue and necessitates creation of a colostomy.⁵³ An APR is necessary in cases where a margin-negative resection of the tumor would result in loss of anal sphincter function resulting in incontinence. Although preoperative chemoRT may result in tumor downsizing and a decrease in tumor bulk (See section on Neoadjuvant/Adjuvant Therapy, below), tumor location is not altered. Whereas sphincter preservation may become possible in cases where initial tumor bulk prevented consideration of such surgery but exposure to the tumor is improved by chemoRT, an APR should be performed when tumor directly involves the anal sphincter or the levator muscles. The lymphatic drainage regions of rectal tumors are influenced by their position in the rectum. More distal tumors are more likely to be characterized by both upward and lateral lymphatic drainage whereas the likelihood of only upward mesorectal drainage is much higher for more proximal tumors.⁵⁴ The TME approach is designed to radically remove lymphatic drainage regions of tumors located above the level of the levator muscles.⁵⁵ The panel does not recommend extension of nodal dissection beyond the field of resection (eg, into the distribution of iliac lymph nodes) unless these nodes are clinically suspicious.

Neoadjuvant/Adjuvant Therapy

Neoadjuvant/adjuvant therapy of rectal cancer often includes locoregional treatment due to the relatively high risk of locoregional recurrence. This risk is associated with the close proximity of the rectum to pelvic structures and organs, the absence of a serosa surrounding the rectum, and technical difficulties associated with obtaining wide surgical margins at resection. In contrast, adjuvant treatment of colon cancer is more focused on preventing distant metastases since this disease is characterized by lower rates of local recurrence.

Combined-modality therapy consisting of surgery, radiation therapy (RT), and chemotherapy is recommended for the majority of patients with stage II (node-negative disease with tumor penetration through the muscle wall) or stage III rectal cancer (node-positive disease without distant metastasis). Use of perioperative pelvic RT in the treatment of patients with stage II/III rectal cancer continues to evolve. Concurrent fluoropyrimidine-based chemotherapy is recommended with radiation.

Ionizing radiation to the pelvis provides local tumoricidal therapy. Putative advantages to preoperative radiation are related to both tumor response and normal tissue.^{56,57} Reducing tumor volume may facilitate resection and increase the likelihood of a sphincter-sparing procedure. Irradiating tissue that is surgery-naïve and thus better oxygenated may result in increased sensitivity to RT. Preoperative radiation can avoid the occurrence of radiation-induced injury to small bowel trapped in the pelvis by post-surgical adhesions. Preoperative radiation that includes structures that will be resected increases the likelihood that an anastomosis with healthy colon can be performed (ie, the anastomosis remains unaffected by the effects of RT because irradiated tissue is resected). One disadvantage of using preoperative RT is the possibility of over-treating early-stage tumors which do not require adjuvant radiation.⁵⁷⁻⁵⁹ Improvements in preoperative staging techniques, such

as endoscopic ultrasound and CT scans, allow for more accurate staging, although the risk of over-staging disease has not been eliminated.⁶⁰

The results of the Swedish Rectal Cancer Trial evaluating the use of short course (5 day) RT administered preoperatively for resectable rectal cancer showed a survival advantage and a decreased rate of local recurrence with this approach compared with surgery alone.⁶¹ However, whereas a number of other studies investigating the effectiveness of preoperative RT or postoperative RT in patients with rectal cancer staged as T1-3 have demonstrated improvements in local control of disease, overall survival was not shown to be significantly affected.^{38,62,63} Preliminary results from a study of patients with stage II/III rectal cancer comparing short course preoperative RT with a postoperative approach which included chemoRT in selected patients (ie, those with a positive CRM following resection) and no RT in patients without evidence of residual disease following surgery indicated that patients in the preoperative RT arm had significantly lower local recurrence rates and a 5% absolute improvement in 3-year disease-free survival (DFS) (P=0.03).⁶⁴ Currently, however, short course RT for the treatment of rectal cancer is not widely practiced in the U.S.

A number of randomized trials have evaluated the effectiveness of chemoRT administered either preoperatively following clinical evaluation/staging (eg, T3-4 by endoscopic ultrasound) or postoperatively following pathologic staging of rectal cancer as T3 and/or N1-2. Putative benefits of addition of chemotherapy concurrent with either pre- or postoperative RT include local RT sensitization and systemic control of disease (ie, eradication of micrometastases), whereas preoperative chemoRT also has the potential to increase rates of pathologic complete response and sphincter preservation. In a study of patients with T3/4 rectal cancer without evidence of distant

metastases who were randomly assigned to receive either preoperative RT alone or preoperative concurrent chemoRT with 5-FU/LV, no difference in overall survival or sphincter preservation was observed in the 2 groups, although patients receiving chemoRT were significantly more likely to exhibit a pathologic complete response (11.4% vs 3.6%; P<0.05) and grade 3/4 toxicity (14.6% vs 2.7%; P<0.05) and less likely to exhibit local recurrence of disease (8.1% vs 16.5%; P<0.05).⁶⁵ A large prospective, randomized trial from The German Rectal Cancer Study Group compared preoperative versus postoperative chemoRT in the treatment of clinical stage II/III rectal cancer.⁵⁷ Results of this study indicated that preoperative therapy was associated with a significant reduction in local recurrence (6% vs 13%; P=0.006) and treatment-associated toxicity, although overall survival was similar in the 2 groups. Preliminary results of a phase III trial that included an evaluation of the addition of chemotherapy to preoperative RT in patients with T3-T4 resectable rectal cancer demonstrated that use of 5-FU/LV chemotherapy enhanced the tumorocidal effect of RT when the 2 approaches were used concurrently. Significant reductions in tumor size, pTN stage, and lymphatic, vascular and perineural invasion rates were observed with use of combined-modality therapy compared with use of RT and surgery without chemotherapy.^{66,67} More mature results from this trial which included 4 treatment groups (preoperative RT; preoperative chemoRT; preoperative RT plus postoperative chemotherapy; and preoperative chemoRT plus postoperative chemotherapy) indicated that no significant differences in overall survival were associated with adding 5-FU-based chemotherapy preoperatively or postoperatively.⁶⁸ Although local recurrence rates were significantly lower in the groups receiving RT followed by chemotherapy, concurrent chemoRT, or concurrent chemoRT plus chemotherapy compared to the group receiving preoperative RT alone, the addition of chemotherapy after concurrent chemoRT did not significantly impact local recurrence rates. In subsequent exploratory analyses of data from the group of patients in this trial who underwent

complete tumor resection without evidence of distant disease before or at surgery, those patients with disease characterized as ypT0-2 showed significant benefit from adjuvant chemotherapy with respect to DFS and overall survival.⁶⁹ These findings may indicate that patients are more likely to benefit from adjuvant therapy if their disease can be downstaged by chemoRT.

Whereas reports from at least one of these studies has indicated that preoperative chemoRT is associated with increased rates of sphincter preservation in rectal cancer patients,⁵⁷ this conclusion has not been supported by 2 recent meta-analyses of randomized trials involving preoperative chemoRT in the treatment of rectal cancer.^{70,71}

Although combined-modality therapy has been associated with decreased rates of local recurrence of rectal cancer, it is also associated with increased toxicity (eg, radiation-induced injury, hematologic toxicities, etc.) relative to surgery alone.^{9,72} It has been suggested that some patients with disease at lower risk of local recurrence (eg, proximal rectal cancer staged as T3, N0, M0, characterized by clear margins and favorable prognostic features) may be adequately treated with surgery and adjuvant chemotherapy.^{9,73,74} Nevertheless, results from a recent retrospective analysis showed the risk of locoregional recurrence to be significantly higher in patients with pT3N0 rectal cancer who did not undergo RT.⁷⁵ In addition, 22% of 188 patients clinically staged with T3N0 rectal cancer by either EUS or MRI who subsequently received preoperative chemoRT had positive lymph nodes following pathologic review of the surgical specimens according to results of a recent retrospective multicenter study.⁶⁰

With respect to the type of chemotherapy administered concurrently with RT, results from the Intergroup 0114 trial, showed bolus 5-FU as part of adjuvant therapy for rectal cancer to be noninferior to bolus 5-FU plus LV.⁷³ After a median follow-up of 4 years, neither the rate of local control nor survival differed among 3 different combinations of

modulated 5-fluorouracil (5-FU) chemotherapy. The equivalence of bolus 5-FU/LV and infusional 5-FU in concurrent chemoRT for rectal cancer is supported by the results of a phase III trial (median follow-up of 5.7 years) in which similar outcomes with respect to overall survival and relapse-free survival were observed when a continuous infusion of 5-FU or bolus 5-FU plus LV was administered concurrently with postoperative RT, although hematologic toxicity was greater in the group of patients receiving bolus 5-FU.⁷⁶ However, results from an earlier trial from the North Central Cancer Treatment Group (NCCTG) showed that postoperative administration of continuous infusion 5-FU during pelvic irradiation was associated with longer overall survival when compared to bolus 5-FU.⁷⁷ Most of the patients in this study had node-positive disease. No phase III randomized data are currently available on the use of capecitabine/RT in rectal cancer, although trials are pending.⁷⁸ A limited number of phase I/II studies have demonstrated that chemoRT with capecitabine was well tolerated with no toxicity or mild to moderate toxicity in the majority of patients with stage II/III rectal cancer and produced results similar to those obtained with continuous infusion of 5-FU and RT.⁷⁹⁻⁸³ A retrospective, matched pair analysis showed no difference in pathologic response, local and distant failure, and overall survival in patients treated with preoperative RT and concurrent capecitabine compared with those treated with preoperative RT and concurrent infusional 5-FU.⁸⁴ Furthermore, results from the study of Smalley et al.⁷⁶ indicating that bolus 5-FU is equivalent to infusional 5-FU in concurrent chemoRT for locally advanced rectal cancer provide indirect support for the hypothesis that capecitabine will not be inferior to 5-FU when used in concurrent chemoRT to treat rectal cancer.

Postoperative chemoRT regimens commonly employ a “sandwich” approach – whereby chemotherapy (typically 5-FU based) is administered before and after the chemoRT regimen.^{73,76,77} The use of FOLFOX or capecitabine chemotherapy before and after postoperative

chemoRT is an extrapolation of the available data in colon cancer.^{85,86} Clinical trials evaluating these agents in the setting of rectal cancer are still pending.⁸⁷

With respect to administration of RT, multiple RT fields should include the tumor or tumor bed with a 2-5 cm margin, presacral nodes, and the internal iliac nodes. The external iliac nodes should also be included for T4 tumors involving anterior structures and the inguinal nodes should be included for tumors invading into the distal anal canal.

Recommended doses of radiation are typically 45-50 Gy, with the exceptions of unresectable cancers where doses higher than 54 Gy may be required, and irradiation of the small bowel where the dose should be limited to 45 Gy. Although not standard routine practice, use of intensity modulated radiotherapy (IMRT) which uses computer-imaging to focus RT to the tumor site and potentially decrease toxicity to normal tissue,^{88,89} can be considered.⁹⁰ As an additional boost, intraoperative radiotherapy (IORT),⁹¹⁻⁹³ which involves direct exposure of tumors to RT during surgery while removing normal structures from the field of treatment should be considered preoperatively for patients with T4 tumors or recurrent cancers to facilitate resection.

Coordination of preoperative therapy, surgery and adjuvant chemotherapy is important. For patients treated with preoperative chemoRT, the panel recommends an interval of 5-10 weeks following completion of full dose 5 ½ week chemoRT prior to performance of surgical resection in order to allow patient recuperation from chemoRT-associated toxicities. Although longer intervals from completion of chemoRT to surgery have been shown to be associated with an increase in pathologic complete response rates,⁹⁴⁻⁹⁶ it is unclear whether this is associated with clinical benefit. Nevertheless, when longer intervals are clinically necessary, they do not appear to increase the blood loss, time associated with surgery, or positive margin rate.⁹⁷

Adjuvant chemotherapy of approximately 4 months duration is recommended for all patients with stage II/III rectal cancer following neoadjuvant chemoRT/surgery regardless of the surgical pathology results, although few studies have evaluated the effect of adjuvant chemotherapy in patients with rectal cancer and its role is not well defined. Evaluation of adjuvant chemotherapy with 5-FU/LV alone versus postoperative RT followed by adjuvant chemotherapy with 5-FU/LV in patients with stage II/III rectal cancer in the National Surgical Breast and Bowel Project (NSABP) R-02 trial showed a significant decrease in local recurrence rate in the group receiving adjuvant chemotherapy after RT compared to the group receiving adjuvant chemotherapy alone.⁹⁸ However, no benefit of adding 5-FU-based adjuvant chemotherapy to preoperative chemoRT with respect to rate of local recurrence was observed in the European Organization for Research and Treatment of Cancer (EORTC) Radiotherapy Group Trial 22921 (hazard ratio=0.87; 95% CI, 0.72-1.04; P=0.13) when the DFS of patients receiving adjuvant chemotherapy following preoperative RT (+/- 5-FU-based chemotherapy) was compared to DFS of patients who underwent preoperative RT (+/- 5-FU-based chemotherapy) but did not receive adjuvant 5-FU-based chemotherapy.⁶⁸ Most of the support for use of FOLFOX or capecitabine as adjuvant chemotherapy in rectal cancer is an extrapolation from the data available for colon cancer.^{85,86} The phase III ECOG E3201 trial is investigating the effect of adding either oxaliplatin (FOLFOX) or irinotecan (FOLFIRI) to 5-FU/LV-based adjuvant chemotherapy administered to stage II/III rectal cancer patients following either preoperative or postoperative chemoRT. Early reports indicate that adjuvant FOLFOX can be safely used in this patient population.⁹⁹ The ECOG E5204 trial is currently evaluating the effect of postoperative 5-FU/LV plus oxaliplatin with or without bevacizumab on the overall survival of patients with stage II/III rectal cancer treated with preoperative 5-FU-based chemoRT. A summary of ongoing clinical trials in early-stage rectal cancer has been presented.⁸⁷

Treatment of Nonmetastatic Rectal Cancer

Recommendations for patients with T1 and T2 lesions

Node-negative T1 and T2 lesions are treated with transabdominal resection or transanal excision (category 2B for T2), if appropriate. This recommendation is category 2B for node-negative T2 tumors since local recurrence rates of 11% to 45% have been observed for T2 lesions following local excision alone.^{39,100,101} In selected lesions that are staged by endoscopic ultrasound or MRI as T1-2, N0 and without adverse pathologic features (eg, no lymphovascular invasion [LVI] or perineural invasion; size less than 3 cm; well to moderately differentiated), local excision with negative margins may give results comparable to transabdominal resection.¹⁰² No additional therapy is recommended for patients with well-differentiated T1 cancers. If pathology review after local excision reveals a poorly differentiated histology, positive margins, or LVI, then a transabdominal re-resection should be performed. T2 cancers excised with negative margins and no poor prognostic factors should be treated with transabdominal resection or adjuvant 5-FU/RT. Systemic chemotherapy should be considered as an adjuvant treatment for those patients who receive adjuvant chemoradiation without additional surgery in order to avoid the risk of undertreatment as the lymph node status is unknown.

For patients with T1 to T2 lesions not amenable to local excision, a transabdominal resection is required. No adjuvant therapy is indicated for patients with pathologic findings of T1 or T2 lesions. Patients with pathologic lymph node-negative T3 lesions (pT3, N0, M0) or pathologic lymph node-positive lesions (pT1-3, N1-2) should receive a “sandwich regimen” consisting of adjuvant chemotherapy with 5-FU with or without LV or FOLFOX (category 2B) or capecitabine (category 2B), followed by concurrent 5-FU/RT (continuous infusion [category 2A] or bolus infusion along with LV [category 2B]) or capecitabine/RT (category 2B), then 5-FU with or without LV or FOLFOX (category 2B) or capecitabine (category 2B). The recommended duration of adjuvant therapy is 6

months. For patients with pathologic evidence of proximal T3, N0, M0 disease with clear margins and favorable prognostic features following an upfront resection, the incremental benefit of RT is likely to be small and chemotherapy alone can be considered, although most patients are not likely to be part of this subset.

Recommendations for patients with T3 lesions and lesions with nodal involvement

Patients clinically staged as having resectable T3, N0 or any T, N1-2 lesions should initially be treated with preoperative combined-modality therapy. Upfront surgery should be reserved for patients with medical contraindications to chemoRT. Preoperative continuous infusional 5-FU/RT is the preferred treatment option (category 1 for node positive disease). Alternative regimens include bolus 5-FU/LV /RT (category 2A) or capecitabine/RT (category 2B). Patients who receive preoperative radiotherapy should undergo transabdominal resection 5-10 weeks following completion of neoadjuvant therapy followed by 6 months of adjuvant chemotherapy (regardless of surgical pathology results) with 5-FU with or without LV (category 1 for T3, N0 or Tany, N1-2 tumors) or FOLFOX (category 2B) or capecitabine (category 2B).

Patients with disease characterized as T3, N0 or T any, N1-2 disease initially treated by transabdominal resection with subsequent pathologic staging of disease as pT1-2, N0, M0 can be followed with observation only. Patients with disease staged as pT3, N0, M0 or pT1-3, N1-2, M0 following initial treatment by transabdominal resection should receive 6 months of adjuvant therapy with 5-FU with or without LV or FOLFOX (category 2B) or capecitabine (category 2B), followed by concurrent 5-FU/RT (5-FU as continuous infusion [category 2A] or bolus infusion with LV [category 2B]) or capecitabine/RT (category 2B), then 5-FU with or without LV (category 2A) or FOLFOX (category 2B) or capecitabine (category 2B). For some patients with pathologic evidence of proximal T3, N0, M0 disease with clear margins and favorable prognostic

features following transabdominal resection, the incremental benefit RT is likely is small and chemotherapy alone can be considered, although this subset of patients is small.

Recommendations for patients with T4 lesions and/or locally unresectable disease

Patients with T4 and/or locally unresectable disease are treated with preoperative continuous infusional 5-FU/RT (category 2A) or bolus 5-FU with LV/RT (category 2A) or capecitabine/RT (category 2B). If possible, resection should be considered following preoperative chemotherapy. Adjuvant therapy for 6 months with either 5-FU with or without LV (category 2A), FOLFOX (category 2B) or capecitabine (category 2B) is indicated regardless of the surgical pathology results.

Treatment of Metastatic Disease

Approximately 50%-60% of patients diagnosed with colorectal cancer will develop colorectal metastases.^{103,104} Patients with stage IV (any T, any N, M1) colorectal cancer or recurrent disease can present with synchronous liver or lung metastases or abdominal peritoneal metastases. Approximately 15%-25% of patients with colorectal cancer present with synchronous liver metastases, although 80%-90% of these patients are initially evaluated to have unresectable metastatic liver disease.^{103,105-107} Metastatic disease more frequently develops metachronously following treatment for colorectal cancer, with the liver as a common site of involvement.¹⁰⁸ There is some evidence to indicate that synchronous metastatic colorectal liver disease is associated with a more disseminated disease state and a worse prognosis than metastatic colorectal disease that develops metachronously. In one retrospective study of 155 patients who underwent hepatic resection for colorectal liver metastases, patients with synchronous liver metastases had more sites of liver involvement (P=0.008) and more bilobar metastases (P=0.016) when compared with patients diagnosed with metachronous liver metastases.¹⁰⁹

It has been estimated that over one-half of patients who die of colorectal cancer have liver metastases at autopsy, and that metastatic liver disease is the cause of death in the majority of these patients.¹¹⁰ Results from reviews of autopsy reports of patients dying from colorectal cancer showed that the liver was the only site of metastatic disease in one-third of patients.¹⁰⁵ Furthermore, rates of 5-year survival for patients with metastatic liver disease not undergoing surgery have been shown to approach 0% in a number of studies.^{103,111} However, studies of selected patients undergoing surgery to remove colorectal liver metastases have demonstrated that cure is possible in this population and should be the goal for many patients with colorectal metastatic liver disease.^{103,112} Recent reports have shown 5-year survival rates following resection of hepatic colorectal metastases exceeding 50%.^{113,114} Therefore, decisions relating to patient suitability, or potential suitability, and subsequent selection for metastatic colorectal surgery are critical junctures in the management of metastatic colorectal liver disease.¹¹⁵

The criteria for determining patient suitability for resection, or surgical cure, of metastatic disease are evolving, with the emphasis being increasingly placed on the likelihood of achieving negative surgical margins while maintaining adequate liver reserve, as opposed to other criteria, such as the number of liver metastases present.¹¹⁶⁻¹¹⁹ Resectability differs fundamentally from endpoints which focus more on palliative measures of treatment such as response and DFS. Instead, the resectability endpoint is focused on the potential of surgery to cure the disease,¹²⁰ since partial liver resection or debulking has not been shown to be beneficial.^{104,118,118} Approaches used in the surgical treatment of liver metastases include preoperative portal vein embolization for the purpose of increasing the volume and function of the portion of the liver which will remain postsurgically, hepatic resection performed in 2 stages for bilobar disease, and the use of ablative methods in combination with resection.¹¹⁶ As with resection,

ablative techniques should be considered only when disease is judged to be completely amenable to ablation.¹²¹ Resection of liver metastases should not be performed in the presence of unresectable sites of extrahepatic disease, and hepatic intra-arterial embolization should not routinely be used outside of a clinical trial. The consensus of the panel is that patients diagnosed with potentially resectable metastatic colorectal cancer should undergo an upfront evaluation by a multidisciplinary team, including surgical consultation (ie, with an experienced hepatic surgeon in cases involving liver metastases) to assess resectability status.

Since the majority of patients diagnosed with metastatic colorectal disease are initially classified as unresectable, preoperative chemotherapy is being increasingly employed to downsize colorectal metastases in order to convert these lesions to a resectable status (ie, conversion chemotherapy); it has also been administered to patients with metastatic disease determined to be resectable (ie, neoadjuvant therapy). Potential advantages of this approach include: earlier treatment of micrometastatic disease; determination of responsiveness to chemotherapy (which can be prognostic and help plan postoperative therapy; and avoidance of local therapy in those who progress early. Potential disadvantages include: chemotherapy-induced liver injury; and missing the “window of opportunity” for resection through the possibility of either disease progression; or achievement of a complete response, thereby making it difficult to identify areas for resection.^{105,122} Furthermore, results from a recent study of colorectal cancer patients receiving preoperative chemotherapy indicated that cancer cells were still present in most of the original sites of metastases when these sites were examined pathologically despite achievement of a complete response as evaluated on CT scan.¹²³ It is therefore essential that during treatment with preoperative chemotherapy, frequent evaluations are undertaken and close communication is maintained between medical oncologists, radiologists, surgeons, and patients so that a

treatment strategy can be developed which optimizes exposure to the preoperative regimen and facilitates an appropriately-timed surgical intervention.¹²⁴ When preoperative therapy is planned, the panel recommends that a surgical re-evaluation should be planned within 8-10 weeks after initiation of preoperative therapy.

Certain clinicopathologic factors, such as the presence of extrahepatic metastases and a disease-free interval of < 12 months, have been associated with a poor prognosis in patients with colorectal cancer,^{113,114,125-127} although the ability of these factors to predict outcome following resection may be limited.¹⁰³ However, decision-making relating to whether to offer preoperative chemotherapy begins with an initial evaluation of the degree of resectability of metastatic disease. Benefits of initial surgery in patients with clearly resectable disease characterized by generally favorable prognostic characteristics may outweigh the benefits of downsizing the disease with neoadjuvant chemotherapy. Alternatively, preoperative chemotherapy would be more appropriate in patients with borderline resectable or initially unresectable but potentially resectable following response to chemotherapy. In addition, preoperative chemotherapy may be more beneficial in patients who have not been exposed to prior chemotherapy or who have not received prior chemotherapy in the previous 12 months.

The most important benefit of the preoperative approach is the potential to convert patients with initially unresectable metastatic disease to a resectable state. In the study of Pozzo et al, it was reported that preoperative therapy with irinotecan combined with 5-FU/LV enabled a significant portion (32.5%) of the patients with initially unresectable liver metastases to undergo liver resection.¹¹⁷ Median time to progression was 14.3 months with all of these patients alive at a median follow-up of 19 months. In a phase II study conducted by the North Central Cancer Treatment Group (NCCTG),¹⁰⁷ 44 patients with unresectable liver

metastases were treated with FOLFOX4. Twenty five patients (60%) had tumor reduction and 17 patients (40%; 68% of the responders) were able to undergo resection after a median period of 6 months of chemotherapy. In another study of 1439 initially unresectable patients with colorectal liver disease, 1104 patients were treated with chemotherapy and 335 patients (23%) were able to undergo primary hepatic resection. Of the 1104 patients receiving chemotherapy, 138 patients (12.5%) classified as “good responders” underwent secondary hepatic resection following preoperative chemotherapy which included oxaliplatin in the majority of cases.¹²⁸ The 5-year survival rate for these 138 patients overall was 33%. More recently, results from a retrospective analysis of 795 previously untreated patients with metastatic colorectal cancer enrolled in the Intergroup N9741 randomized phase III trial evaluating the efficacy of mostly oxaliplatin-containing chemotherapy regimens indicated that 24 patients (3.3%) were able to undergo curative liver resection following treatment.¹²⁹ The median overall survival time in this group was 42.4 months.

Recently, the efficacy of bevacizumab in combination with FOLFOX and FOLFIRI (infusional 5-FU, LV, irinotecan) in the treatment of unresectable metastatic disease (see section on [Chemotherapy for Advanced or Metastatic Disease](#)) has led to its use in combination with these regimens in the preoperative setting, although the safety of administering bevacizumab pre- or postoperatively, in combination with 5-FU-based regimens has not been adequately evaluated. A retrospective evaluation of data from 2 randomized trials of 1132 patients receiving chemotherapy with or without bevacizumab as initial therapy for metastatic colorectal cancer indicated that the incidence of wound healing complications was increased for the group of patients undergoing a major surgical procedure while receiving a bevacizumab-containing regimen when this population was compared to the group receiving chemotherapy alone while undergoing major surgery (13% vs 3.4%, respectively; P=0.28).¹³⁰ However, when chemotherapy plus

bevacizumab or chemotherapy alone was administered prior to surgery, the incidence of wound healing complications in either group of patients was low (1.3% vs 0.5%; P=0.63). The panel recommends at least a 6 week interval (which corresponds to 2 half-lives of the drug¹³¹) between the last dose of bevacizumab and elective surgery. Further support for this recommendation comes from results of a single center, nonrandomized phase II trial of patients with potentially resectable liver metastases which showed no increase in bleeding or wound complications when the bevacizumab component of CapeOX plus bevacizumab therapy was stopped 5 weeks prior to surgery (ie, bevacizumab excluded from the 6th cycle of therapy).¹³² In addition, no significant differences in bleeding, wound, or hepatic complications were observed in a retrospective trial evaluating effects of preoperative bevacizumab stopped ≤ 8 weeks vs. > 8 weeks prior to resection of liver colorectal metastases for patients receiving oxaliplatin- or irinotecan-containing regimens.¹³³

Other reported risks associated with the preoperative approach include the potential for development of liver steatosis or steatohepatitis when oxaliplatin or irinotecan-containing chemotherapeutic regimens are administered.¹²⁴ To limit the development of hepatotoxicity, it is therefore recommended that surgery should be performed as soon as possible after the patient becomes resectable and usually not more than 3-4 months following initiation of preoperative treatment.

Colorectal metastatic disease can also occur in the lung.¹³⁴ Most of the treatment recommendations discussed for metastatic colorectal liver disease, with the exception of hepatic arterial infusion (HAI), also apply to the treatment of colorectal pulmonary metastases. Combined pulmonary and hepatic resections of resectable metastatic disease have been performed in selected cases.¹³⁵ The goal of treatment of most abdominal/peritoneal metastases is palliative, rather than curative.

It is important to note that some of the treatment approaches for patients diagnosed with rectal cancer and potentially resectable synchronous lung or liver metastases differ relative to those for patients diagnosed with stage IV colon cancer characterized as potentially resectable metastatic disease. In particular, initial treatment options for potentially resectable rectal cancer include: preoperative chemoRT directed toward treatment of the primary cancer; preoperative combination chemotherapy with a bevacizumab-containing regimen to target metastatic disease; and a surgical approach (ie, staged or synchronous resection of metastases and rectal lesion). Advantages of an initial chemoRT approach include a possible decreased risk of pelvic failure following surgery although preoperative pelvic RT may decrease tolerance to systemic bevacizumab-containing adjuvant regimens, thereby limiting subsequent treatment of systemic disease. However, data to guide decisions regarding optimal treatment approaches in this population of patients are very limited. Of note, patients with stage II/III rectal cancer enrolled in a large randomized trial evaluating the effect of adding chemotherapy to preoperative RT were found to be three times more likely to develop distant metastases than local recurrence of disease after a median follow-up of over 5 years.⁶⁸

Only limited data exist regarding the efficacy of adjuvant chemotherapy following resection for metastatic colorectal liver or lung disease. Nevertheless, the panel recommends administration of a course of an active systemic chemotherapy regimen for metastatic disease for some patients following liver or lung resection who have received preoperative chemoRT or no preoperative therapy following staged or synchronous resection of metastases and rectal lesion in order to increase the likelihood that residual microscopic disease will be eradicated. Postoperative chemoRT is recommended for patients with synchronous metastases who have not received prior chemoRT and who are at higher risk for pelvic recurrence following staged or

synchronous resection of metastases and rectal lesion (ie, patients with disease staged as pT3-4, Any N, or Any T,N1-2).

Placement of a hepatic arterial port or implantable pump during surgical intervention for liver resection with subsequent administration of chemotherapy directed to the liver metastases through the hepatic artery (i.e. HAI) is listed in the guidelines as an option (category 2B). In a randomized study of patients who had undergone hepatic resection, administration of floxuridine (with dexamethasone and with or without LV) by HAI in addition to systemic chemotherapy was shown to be superior to systemic chemotherapy alone with respect to 2-year survival and time to progression of hepatic disease.^{136,137} However, the difference in survival between the 2 arms of the study was not significant at later follow-up periods.^{136,138} A number of other clinical trials have shown significant improvement in response or time to hepatic disease progression when HAI therapy was compared with systemic chemotherapy, although most have not shown a survival benefit of HAI therapy.¹³⁶ Some of the uncertainties regarding patient selection for preoperative chemotherapy are also relevant to the application of HAI.¹¹² Limitations on the use of HAI therapy include the potential for biliary toxicity,¹³⁶ and the requirement for specific technical expertise. The consensus of the panel is that HAI therapy should be considered only at institutions with extensive experience in both the surgical and medical oncologic aspects of the procedure.

Although the benefit of preoperative or postoperative chemotherapy for patients with liver metastases has not yet been validated in randomized clinical trials, a recent European Organization for Research and Treatment of Cancer (EORTC) phase III study evaluating use of perioperative FOLFOX4 (6 cycles before and 6 cycles after surgery) for patients with initially resectable liver metastases demonstrated absolute improvements in 3-year PFS of 8.1% (P=0.041) and 9.2% (P=0.025) for all eligible patients and all resected patients, respectively, when

chemotherapy in conjunction with surgery was compared with surgery alone.¹³⁹

Locally recurrent rectal cancer is characterized by isolated pelvic/anastomotic recurrence of disease. In a single-center study at M D Anderson, rates of 5-year local recurrence were reported to be low (ie, 5-year locoregional control rate of 91%) for patients with rectal cancer treated with surgery and either RT or chemoRT, and 78% of recurrences occurred in the low pelvic and presacral regions.¹⁴⁰

Patients with disease recurrence at the anastomotic site are more likely than those with an isolated pelvic recurrence to be cured following re-resection.^{141,142} In a study of 43 consecutive patients with advanced pelvic recurrence of colorectal cancer who had not undergone prior RT, treatment with 5 weeks of 5-FU by continuous infusion concurrent with RT enabled the majority of patients (77%) to undergo re-resection with curative intent.¹⁴¹

Recommendations for Treatment of Synchronous Metastases/Resectable Disease

As part of the pre-treatment work-up, the panel recommends tumor KRAS gene status testing for all patients with metastatic colorectal cancer at the time of diagnosis of metastatic disease (see discussion of KRAS testing on MS-21).

Initial treatment options for patients with stage IV disease (any T, any N, M1) with resectable liver or lung metastases include: staged or synchronous resection of metastases and rectal lesion; treatment with continuous infusional 5-FU/pelvic RT (category 2A) or bolus 5-FU with LV/pelvic RT (category 2A) or capecitabine/RT (category 2B); or combination chemotherapy (eg, FOLFOX, CapeOX, or FOLFIRI regimens with or without bevacizumab). For the latter 2 groups of patients, surgery should be performed 5-10 weeks following completion of neoadjuvant therapy.

Adjuvant therapy for patients undergoing initial surgery is dependent on pathologic staging of disease. For patients undergoing initial surgical treatment, the panel recommends that those at higher risk for pelvic failure relative to systemic disease (eg, disease pathologically staged as pT3-4, Any N or Any T, N1-2) undergo postoperative chemoRT using the “sandwich” approach (ie, chemotherapy followed by concurrent chemoRT followed by chemotherapy for 4-6 months).^{76,77} The panel acknowledged that not all patients with rectal cancer and resectable liver or lung metastases need to be treated with chemoRT. For example, in the population of patients with pT1-2,N0 disease, the competing risk of distant metastases is considered to be higher than that of locoregional recurrence. Therefore, the panel recommended that these patients receive adjuvant chemotherapy with one of the following options: 5-FU with or without LV for 6 months (category 2A); FOLFOX or CapeOX plus bevacizumab for 4-6 months (category 2B); FOLFIRI plus bevacizumab for 4-6 months (category 2B). Adjuvant therapy recommendations for patients who have received neoadjuvant chemoRT is as described for patients with pT1-2,N0 disease, whereas patients who have undergone neoadjuvant bevacizumab-containing therapy should receive postoperative chemoRT as described above for patients with pT3-4, Any N, or Any T, N1-2 disease.

Recommendations for Treatment of Synchronous Metastases/Unresectable Disease

As part of the pre-treatment work-up, the panel recommends tumor KRAS gene status testing for all patients with metastatic colorectal cancer at the time of diagnosis of metastatic disease (see discussion of KRAS testing on MS-21).

Patients with any unresectable or medically inoperable metastases are treated according to whether they are symptomatic or asymptomatic. Symptomatic patients are treated with chemotherapy alone or combined modality therapy with 5-FU/RT or capecitabine/RT (category

2B), resection of the involved rectal segment or laser canalization or diverting colostomy or stenting.

Recommendations for Treatment of Metachronous Metastases

Upon documentation of metachronous metastases in which disease is or may become potentially resectable, characterization of the extent of disease by PET scan is recommended. PET is used at this juncture to promptly characterize the extent of metastatic disease, and to identify possible sites of extrahepatic disease which could preclude surgery.¹⁴³

As with other first identifications of metastatic disease, a tumor sample (metastases or original primary) should be sent for KRAS genotyping in order to define whether anti-EGFR agents can be considered in the list of potential options for this patient (see discussion of KRAS testing on MS-21).

The management of metachronous metastatic disease is further distinguished from that of synchronous disease by also including an evaluation of the chemotherapy history of the patient, and by the absence of transabdominal resection. Resectable patients are classified according to whether they have received no previous chemotherapy or prior chemotherapy within or prior to the previous 12 months. For patients who have not received prior chemotherapy and who have resectable metastatic disease, primary treatment options include initial resection followed by chemotherapy or neoadjuvant chemotherapy followed by resection and additional postoperative chemotherapy; The optimal sequence of therapeutic interventions is less clear for patients who have received prior adjuvant chemotherapy. For patients who exhibit disease recurrence or progression during or within 12 months of chemotherapy, the role of neoadjuvant chemotherapy is less clear. Following surgery, adjuvant therapy with an alternative active metastatic chemotherapy regimen can be considered.

Patients determined by cross-sectional imaging or PET scan to have unresectable rectal cancer should receive an active metastatic

chemotherapy regimen based on prior chemotherapy history. Specifically, patients exhibiting disease progression on FOLFOX administered within the previous 12 months should be switched to a FOLFIRI regimen with the option of inclusion of bevacizumab. Patients with chemotherapy-responsive disease who are converted to a resectable stage should undergo resection followed by adjuvant treatment with an active chemotherapy regimen. If metastatic lesions remain unresectable subsequent treatment is dependent, in part, on the performance status (PS) of the patient. Treatment with an active chemotherapy regimen for advanced or metastatic disease is the treatment of choice for patients with PS 0-2. Patients with PS ≥ 3 are given best supportive care. Best supportive care is an option for patients diagnosed with metachronous metastases who have previously received all active chemotherapy regimens in cases of both resectable and unresectable disease.

Isolated pelvic/anastomotic recurrence is optimally managed by preoperative RT and concurrent infusional 5-FU, if full course RT was not given previously. If full course RT was given previously, additional RT should be considered if it can be safely delivered.¹⁴⁴⁻¹⁴⁶ Resection should be performed, if possible, although debulking, resulting in gross residual cancer, is discouraged. Patients with unresectable lesions are treated according to their ability to tolerate therapy. The goal of treatment for most abdominal/peritoneal metastases is palliative, rather than curative. The panel currently considers the treatment of disseminated carcinomatosis with cytoreductive surgery (ie, peritoneal stripping surgery) and perioperative hyperthermic intraperitoneal chemotherapy.^{147,148} to be investigational and does not endorse such therapy outside of a clinical trial. However, the panel recognizes the need for randomized clinical trials that will address the risks and benefits associated with each of these modalities.

Chemotherapy for Advanced or Metastatic Disease

The current management of disseminated metastatic colon cancer uses various active drugs, either in combination or as single agents: 5-FU/LV, capecitabine; irinotecan, oxaliplatin, bevacizumab, cetuximab, and panitumumab.¹⁴⁹⁻¹⁶⁴ The putative mechanisms of action of these agents are varied and include interference with DNA replication, and inhibition of the activities of vascular endothelial growth factor (VEGF) and epidermal growth factor (EGF) receptors.¹⁶⁵⁻¹⁶⁸ The choice of therapy is based on consideration of the type and timing of the prior therapy that has been administered and the differing toxicity profiles of the constituent drugs. Although the specific chemotherapy regimens listed in the guideline are designated according to whether they pertain to initial therapy, therapy after first progression, or therapy after second progression, it is important to clarify that these recommendations represent a continuum of care and that these lines of treatment are blurred rather than discrete.¹⁵¹ For example, if oxaliplatin, administered as a part of an initial treatment regimen, is discontinued after 12 weeks or earlier for escalating neurotoxicity, continuation of the rest of the treatment regimen would still be considered initial therapy. Principles to consider at the start of therapy include pre-planned strategies for altering therapy for patients in both the presence and absence of disease progression, as well as plans for adjusting therapy for patients who experience certain toxicities. For example, decisions related to therapeutic choices following first progression of disease should be based, in part, on the prior therapies received by the patient (ie, exposing patient to a range of cytotoxic agents). Further, an evaluation of the efficacy and safety of these regimens for an individual patient must take into account not only the component drugs, but also the doses, schedules, and methods of administration of these agents, as well as the potential for surgical cure and the performance status of the patient.

As initial therapy for metastatic disease in a patient with good tolerance to intensive therapy, the panel recommends a choice of 4 chemotherapy regimens: FOLFOX (eg, FOLFOX4 or mFOLFOX6),^{152, 160, 169-175} CapeOX,¹⁷⁵⁻¹⁷⁷ FOLFIRI,^{153,170,174,178} or 5-FU/LV.^{155, 159, 178-180}

The panel further recommends that each of these regimens be administered in combination with bevacizumab when used for initial therapy. With respect to the treatment of metastatic disease, the consensus of the panel was that FOLFOX plus bevacizumab and CapeOX plus bevacizumab can be used interchangeably,¹⁷⁵ and that both of these combination regimens, as well as FOLFIRI plus bevacizumab, represent appropriate standard practices for the initial treatment of metastatic colorectal cancer. The infusional 5-FU/LV plus bevacizumab regimen is recommended as initial therapy for patients not able to tolerate oxaliplatin or irinotecan since it has been shown to be associated with lower toxicity.¹⁸¹⁻¹⁸⁴

Pooled results from several randomized phase II studies have demonstrated that addition of bevacizumab to first-line 5-FU/LV regimens improved overall survival in patients with metastatic colorectal cancer when compared to survival results for patients receiving these regimens without bevacizumab.^{182, 185} A combined analysis of the results of several of these trials showed that addition of bevacizumab to 5-FU/LV-containing regimens was associated with a median survival of 17.9 months versus 14.6 months for regimens consisting of 5-FU/LV or 5-FU/LV plus irinotecan without bevacizumab.¹⁸⁵ A study of previously untreated patients receiving bevacizumab and irinotecan-5-FU chemotherapy (IFL) also provided support for the inclusion of bevacizumab in initial therapy.¹⁸⁴ In that pivotal trial a markedly longer survival time was observed with the use of bevacizumab: 20.3 months versus 15.6 months (hazard ratio for death = 0.66; P<0.001). Results from a recent head-to-head randomized, double-blind, placebo-controlled phase III study (N016966) comparing CapeOX (capecitabine dose 1000 mg/m² twice daily for 14 days) with FOLFOX have been

reported. With a median follow-up period of over 30 months, results from this study support the conclusion that neither regimen is inferior with respect to the other in terms of toxicity or efficacy endpoints when used in the initial treatment of metastatic colorectal cancer.^{175, 186-188} In this trial, addition of bevacizumab to oxaliplatin-based regimens was associated with an increase in progression-free survival (PFS) compared to these regimens without bevacizumab (hazard ratio=0.83; 97.5% CI, 0.72-0.95; P=0.0023). However, the significant incremental benefit observed with addition of bevacizumab was more modest than seen in some earlier trials, and it has been suggested that differences observed in cross-study comparisons of N016966 with other trials might be related to differences in the discontinuation rates and durations of treatment between trials,¹⁸⁹ although such hypotheses are only conjectural. Furthermore, in this 1400 patient randomized study, absolutely no difference in response rates was seen with and without bevacizumab (see below), and this finding would not be potentially influenced by the early withdrawal rates, which occurred after the responses would have occurred. Results of subset analyses evaluating the benefit of adding bevacizumab to either FOLFOX or CapeOX indicated that bevacizumab was associated with improvements in PFS when added to CapeOX but not FOLFOX, although the PFS curves observed for patients receiving either CapeOX plus bevacizumab or FOLFOX plus bevacizumab were nearly identical.¹⁸⁶ An analysis of the ITT population demonstrated no statistically significant increase in median overall survival for patients in the bevacizumab-containing arm of the N016966 trial (21.3 vs. 19 months) (hazard ratio=0.89; 97.5% CI, 0.76-1.03; P=0.0769).¹⁸⁸ The results of the phase III BICC-C study evaluating the effectiveness of 3 irinotecan-containing regimens with and without bevacizumab demonstrated that, for first-line treatment of advanced colorectal cancer, FOLFIRI is superior to a modified IFL regimen or CapeIRI (capecitabine plus irinotecan) in terms of efficacy and safety.^{190, 191} Although this trial was closed early and did not meet projected enrollment, a significant increase in PFS was observed for

patients receiving first-line FOLFIRI (7.6 months) when compared to PFS results for patients receiving either a modified IFL regimen (5.9 months; P=0.004) or CapeIRI (5.8 months; P=0.015) at a median follow-up of 22.6 months, although no significant differences in median overall survival were observed for the modified IFL or CapeIRI regimens compared with the FOLFIRI regimen. When FOLFIRI or modified IFL was combined with bevacizumab, PFS was shown to increase to 11.2 and 8.3 months, respectively, although this difference was not significant (P=0.28). However, at a median follow-up of 34.4 months, overall survival was significantly higher for patients receiving FOLFIRI plus bevacizumab (28.0 months) compared with modified IFL plus bevacizumab (19.2 months; P=0.037).¹⁹¹ Evidence for the comparable efficacy for FOLFOX and FOLFIRI comes from a crossover study in which patients received either FOLFOX or FOLFIRI as initial therapy and were then switched to the other regimen at the time of disease progression.¹⁷⁰ Similar response rates and PFS times were obtained when these 2 regimens were used as first-line therapy. Further support for this conclusion has come from results of a phase III trial comparing the efficacy and toxicity of FOLFOX4 and FOLFIRI regimens in previously untreated patients with metastatic colorectal cancer.¹⁷⁴ No differences were observed in response rate, PFS times, and overall survival in the 2 treatment arms. The results of an ongoing phase III study evaluating the effectiveness of FOLFIRI in combination with bevacizumab in the initial treatment of patients with metastatic disease have not yet been reported.¹⁹²

Convincing, albeit indirect, support for inclusion of bevacizumab in combination with chemotherapeutic agents in the initial treatment of advanced or metastatic colon cancer comes from results of the randomized phase III study E3200, conducted by Eastern Cooperative Oncology Group (ECOG), which demonstrated that bevacizumab in combination with FOLFOX4 improved survival in bevacizumab-naïve patients with previously-treated advanced colorectal cancer. Median

overall survival was 12.9 months for patients receiving FOLFOX4 plus bevacizumab compared to 10.8 months for patients receiving FOLFOX4 alone (P=0.0011).¹⁹³ Use of single agent bevacizumab is not recommended since it was shown to have inferior efficacy compared with the FOLFOX alone or FOLFOX plus bevacizumab treatment arms.¹⁹³ Although this study involved patients with previously-treated disease, the results cannot be used to support use of bevacizumab in patients after first or second progression if they have progressed on a bevacizumab-containing regimen.

The risk of stroke and other arterial events is increased in elderly patients receiving bevacizumab.¹⁹⁴ In addition, use of bevacizumab may interfere with wound healing^{183,194,195} (see [Treatment of Metastatic Disease](#)), and gastrointestinal perforation is a relatively rare, but important, side effect of bevacizumab therapy in patients with colorectal cancer.^{183,195} Extensive prior intra-abdominal surgery, such as peritoneal stripping, may predispose patients to gastrointestinal perforation. A small cohort of patients with advanced ovarian cancer had an unacceptably high rate of gastrointestinal perforation when treated with bevacizumab¹⁹⁶; this illustrates that peritoneal debulking surgery may be a risk factor for gastrointestinal perforation whereas the presence of an intact primary tumor does not appear to increase risk of gastrointestinal perforation.

With respect to the toxicities associated with capecitabine use, the panel noted that patients with diminished creatinine clearance may accumulate levels of the drug,¹⁹⁷ that the incidence of hand-foot syndrome was increased for patients receiving capecitabine-containing regimens versus either bolus or infusional regimens of 5-FU/LV^{183, 197} and that North American patients may experience a higher incidence of adverse events with certain doses of capecitabine compared with patients from other countries.¹⁹⁸ Such toxicities may necessitate modifications in the dosing of capecitabine,^{183, 197, 199} and patients on

capecitabine should be monitored closely so that dose adjustments can be made at the earliest signs of certain side effects such as hand-foot syndrome. It is currently not known whether the efficacy of CapeOX plus bevacizumab and FOLFOX plus bevacizumab remain comparable when capecitabine doses are lower than the 1000 mg/m² twice daily dose used in the study of Saltz et al.¹⁸⁶

Toxicities associated with irinotecan include both early and late forms of diarrhea, dehydration, and severe neutropenia.^{200, 201} Irinotecan is metabolized by the enzyme uridine diphosphate-glucuronyl transferase 1A1 (UGT1A1) which is also involved in converting substrates, such as bilirubin, into more soluble forms through conjugation with certain glycosyl groups. Deficiencies in UGT1A1 can be caused by certain genetic polymorphisms, and can result in conditions associated with accumulation of unconjugated hyperbilirubinemias, such as types I and II of Crigler-Najjar syndrome and Gilbert syndrome. Thus, irinotecan should be used with caution and at decreased dose in patients with Gilbert's disease or elevated serum bilirubin.²⁰² Similarly, certain genetic polymorphisms in the gene encoding for UGT1A1 can result in a decreased level of glucuronidation of the active metabolite of irinotecan, resulting in an accumulation of the drug,^{201, 203} although severe irinotecan-related toxicity is not experienced by all patients with these polymorphisms.²⁰³ A commercial test is available to detect the UGT1A1*28 allele which is associated with decreased gene expression and, hence, reduced levels of UGT1A1 expression,²⁰² and a new warning has been added to the label for Camptosar which indicates that a reduced starting dose of the drug should be used in patients known to be homozygous for UGT1A1*28.²⁰⁰ A practical approach to the use of UGT1A1*28 allele testing with respect to patients receiving irinotecan has been presented,²⁰³ although guidelines for the use of this test in clinical practice have not been established. Furthermore, UGT1A1 testing on a patient who has experienced irinotecan toxicity is not recommended since that patient will require a dose reduction

regardless of the UGT1A1 test result. Use of oxaliplatin has been associated with an increased incidence of peripheral sensory neuropathy.²⁰⁴ Results of the OPTIMOX1 study demonstrated that a “stop-and-go” approach employing oxaliplatin-free intervals resulted in decreased neurotoxicity but did not affect overall survival in patients receiving FOLFOX as initial therapy for metastatic disease. Therefore, the panel recommends adjustments in the schedule/timing of the administration of this drug as a means of limiting this adverse effect.²⁰⁵ Discontinuation of oxaliplatin from FOLFOX or CapeOX should be strongly considered after 3 months of therapy, or sooner for unacceptable neurotoxicity) with other drugs in the regimen maintained until time of tumor progression. Patients experiencing neurotoxicity on oxaliplatin should not receive subsequent oxaliplatin therapy until and unless there is near-total resolution of that neurotoxicity, but oxaliplatin can subsequently be reintroduced if stopped to prevent development of neurotoxicity. In the phase II OPTIMOX2 trial, patients were randomized to receive an induction FOLFOX regimen (6 cycles) followed by discontinuation of all chemotherapy until tumor progression reached baseline followed by reintroduction of FOLFOX or an OPTIMOX1 approach (discontinuation of oxaliplatin after 6 cycles of FOLFOX [to prevent or reduce neurotoxicity] with continuance of 5-FU/LV followed by reintroduction of oxaliplatin upon disease progression).²⁰⁶ Results of the study demonstrated a strong trend for improved overall survival for patients receiving the OPTIMOX1 approach compared with patients undergoing an early, pre-planned chemotherapy-free interval (median overall survival 26 vs. 19 months; $P=0.0549$).

The consensus of the panel is that infusional 5-FU regimens appear to be less toxic than bolus regimens and that any bolus regimen of 5-FU is inappropriate when administered with either irinotecan or oxaliplatin. Therefore, the panel no longer recommends using the IFL (irinotecan, bolus 5-FU/LV) regimen (which was shown to be associated with

increased mortality and decreased efficacy relative to FOLFIRI in the BICC-C trial¹⁹⁰ and inferior to FOLFOX in the Intergroup trial¹⁵²) at any point in the therapy continuum and it has been removed from the guidelines. 5-FU in combination with irinotecan or oxaliplatin should be administered via an infusional biweekly regimen,^{159,178} or capecitabine should be used.¹⁵⁶

The recommended therapy options after first progression for patients who have received prior 5-FU/LV-based therapy are dependent on the initial treatment regimen and include FOLFIRI¹⁷⁸ with or without cetuximab, and irinotecan in combination with cetuximab¹⁶² or as a single agent,¹⁵⁴ for patients who had received a FOLFOX or CapeOX-based regimen for initial therapy. FOLFOX or CapeOX alone is an option for patients who received a FOLFIRI-based regimen as initial treatment. The recommendations regarding use of CapeOX in lieu of FOLFOX after first progression are supported by the results of studies demonstrating comparable efficacy of these 2 agents in initial therapy.¹⁷⁵ Other options for patients initially treated with a FOLFIRI-based regimen include cetuximab plus irinotecan, or single agent cetuximab or panitumumab for those not able to tolerate the combination with irinotecan. For patients receiving 5-FU/LV without oxaliplatin or irinotecan as initial therapy, options after first progression include: FOLFOX, CapeOX, FOLFIRI or single agent irinotecan.

Results from a randomized study to evaluate the efficacy of FOLFIRI and FOLFOX6 regimens as initial therapy and to determine the effect of using sequential therapy with the alternate regimen following first progression showed neither sequence to be significantly superior with respect to PFS or median overall survival.¹⁷⁰ A combined analysis of data from 7 recent phase III clinical trials in advanced colorectal cancer provided support for a correlation between an increase in median survival and administration of all of the 3 cytotoxic agents (ie, 5-FU/LV, oxaliplatin, and irinotecan) at some point in the continuum of care.²⁰⁷

Furthermore, overall survival was not found to be associated with the order in which these drugs were received. Single agent irinotecan administered after first progression has been shown to significantly improve overall survival relative to best supportive care²⁰⁸ or infusional 5-FU/LV.²⁰⁹ In the study of Rougier et al.,²⁰⁹ median overall survival was 4.2 months for irinotecan versus 2.9 months for 5-FU (P=0.030) whereas Cunningham et al.²⁰⁸ reported a survival rate at 1 year of 36.2% in the group receiving irinotecan versus 13.8% in the supportive-care group (P=0.001). Furthermore, no significant differences in overall survival were observed in the Intergroup N9841 trial when FOLFOX was compared to irinotecan monotherapy following first progression of metastatic colorectal cancer.²¹⁰ Infusion of calcium and magnesium salts has been suggested as a potential means of limiting the neurotoxic effects of oxaliplatin. Data are limited on this topic but such an approach may be considered.

A sizable body of literature has demonstrated that the status of the KRAS gene in the tumor is highly predictive of outcome with anti-EGFR therapies.²¹¹⁻²²⁰ Tumors that have a mutation in codon 12 or codon 13 of the KRAS gene are essentially insensitive to EGFR inhibitors such as cetuximab or panitumumab. The panel therefore strongly recommends KRAS genotyping of tumor tissue (either primary tumor or metastasis) in all patients with metastatic colorectal cancer. Patients with known codon 12 or 13 KRAS mutations should *not* be treated with either cetuximab or panitumumab, either alone or in combination with other anticancer agents, as there is virtually no chance of benefit and the exposure to toxicity and expense cannot be justified. It is to be emphasized that KRAS mutations are early events in colorectal cancer formation, and there is a tight correlation between mutation status in the primary tumor and the metastases.²²¹ For this reason, KRAS genotyping can be done on archived specimens of either the primary tumor or a metastasis. Fresh biopsies should not be obtained solely for

the purpose of KRAS genotyping if an archived specimen from either the primary or a metastasis is available.

Cetuximab has been studied trial as both a single agent^{162, 222} and in combination with irinotecan^{162, 223} in the treatment of metastatic colorectal cancer. A partial response rate of 9% was observed when single agent cetuximab was administered in an open-label phase II trial to 57 patients with colorectal cancer refractory to prior irinotecan-containing therapy.²²² More recently, cetuximab monotherapy was reported to significantly increase both PFS (hazard ratio=0.68; 95% CI, 0.57-0.80; P<0.001) and overall survival (hazard ratio=0.77; 95% CI, 0.64-0.92; P=0.005) for patients with refractory metastatic colorectal cancer when compared with best supportive care alone.²²⁴ Results from a direct comparison of cetuximab monotherapy and the combination regimen of cetuximab and irinotecan in patients who had progressed following initial therapy with an irinotecan-based regimen indicated that response rates were doubled in the group receiving the combination of cetuximab plus irinotecan when compared with patients receiving cetuximab monotherapy (22.9% versus 10.8% [P=0.007]).¹⁶² Results of a large phase III study of similar design did not demonstrate a difference in overall survival between the 2 treatment arms but also showed significant improvement in response rate, and in median PFS, with the combination of irinotecan and cetuximab compared with irinotecan alone. Toxicity was higher in the cetuximab-containing arm.²²⁵ Therefore it is acceptable to use either irinotecan alone or cetuximab plus irinotecan. For patients receiving irinotecan alone, the combination of cetuximab and irinotecan is preferable to cetuximab alone as therapy after progression on irinotecan for those who can tolerate this combination. For patients not able to tolerate cetuximab plus irinotecan, either single agent cetuximab or single agent panitumumab can be considered.

Panitumumab has been studied as a single agent in the setting of metastatic colorectal cancer for patients with disease progression on both oxaliplatin and irinotecan-based chemotherapy¹⁶¹; respective response rates of 10% versus 0% ($P < 0.0001$) for panitumumab plus best supportive care versus best supportive care alone were observed, as well as a significant increase in PFS with panitumumab (hazard ratio=0.54; 95% CI, 0.44-0.66). Results of the PACCE trial showed decreased PFS and increased toxicity of chemotherapy/bevacizumab/panitumumab over chemotherapy/bevacizumab.²²⁶ Thus, recommendations for the use of panitumumab in the guidelines are currently restricted to single agent use only. The panel allows that panitumumab can be substituted for cetuximab when either drug is used as a single agent following first or second progression. Although no head-to-head studies comparing cetuximab and panitumumab have been undertaken, this recommendation is supported by the similar response rates observed when each agent was studied as monotherapy. One difference between these 2 agents is that panitumumab is a fully human monoclonal antibody whereas cetuximab is a chimeric monoclonal antibody.^{227, 228} There are no data to support use of either cetuximab or panitumumab after failure of the other drug and the panel recommends against this practice. Cetuximab in combination with irinotecan is also indicated following progression for patients refractory to irinotecan-based chemotherapy since it has shown activity in this setting.¹⁶² Administration of either cetuximab or panitumumab has been associated with severe infusion reactions, including anaphylaxis, in 3% and 1% of patients, respectively.^{227, 228} Based on case reports, for those patients experiencing severe infusion reactions to cetuximab, administration of panitumumab appears to be feasible.^{229,230} Skin toxicity is a side effect of both of these agents and is *not* considered to be part of the infusion reactions. The incidence and severity of skin reactions with cetuximab and panitumumab appears to be very similar; however, the presence and severity of skin rash in patients receiving

either of these drugs has been shown to be predictive of increased response and survival.^{224,231,232}

EGFR testing of colorectal tumor cells has no demonstrated predictive value in determining likelihood of response to either cetuximab or panitumumab. Data from the BOND study indicated that the intensity of immunohistochemical staining of colorectal tumor cells did not correlate with the response rate to cetuximab.¹⁶² A similar conclusion was drawn with respect to panitumumab.²³³ Therefore, routine EGFR testing is not recommended, and no patient should be either considered for or excluded from cetuximab or panitumumab therapy on the basis of EGFR test results.

With respect to the treatment continuum for metastatic colorectal cancer, there are no data to support the addition of bevacizumab to a regimen following clinical failure of a previous bevacizumab-containing regimen.¹⁹³ Therefore, routine use of cetuximab plus bevacizumab in patients who have experienced clinical failure on a bevacizumab-containing regimen is not recommended.

A recent study of 6,286 patients from 9 trials which evaluated the benefits and risks associated with intensive first-line treatment in the setting of metastatic colorectal cancer treatment according to patient performance status showed similar therapeutic efficacy for patients with performance status=2 or ≤ 1 as compared with control groups, although the risks of certain gastrointestinal toxicities were significantly increased for patients with performance status=2.²³⁴ For patients with impaired tolerance to aggressive initial therapy, the guideline includes recommendations for single-agent capecitabine,^{156,157} or infusional 5-FU/leucovorin,^{158,159} with or without bevacizumab (category 2B for combination with bevacizumab). Although a comparison of capecitabine plus bevacizumab versus capecitabine alone as initial therapy for metastatic cancer has not been done, CapeOX plus bevacizumab has been shown to be superior to CapeOX alone in this setting.^{183,186}

Metastatic cancer patients with no improvement in functional status should receive best supportive care. Patients showing improvement in functional status should be treated with one of the options specified for therapy after first progression as described above. The panel recommends that progression of disease following treatment with an EGFR inhibitor alone or a regimen including cetuximab and irinotecan should be followed by either best supportive care or enrollment in a clinical trial. The panel recommends against the use of capecitabine, mitomycin, alfa-interferon, taxanes, methotrexate, pemetrexed, sunitinib, sorafenib, erlotinib, or gemcitabine, either as single agents or in combination, as salvage therapy in patients exhibiting disease progression following treatment with standard therapies. These agents have not been shown to be effective in this setting. No objective responses were observed when single agent capecitabine was administered in a phase II study of patients with colorectal cancer resistant to 5-FU.²³⁵

Post-Treatment Surveillance

The approach to monitoring and surveillance of patients with rectal cancer is similar to that described for colon cancer with the addition of proctoscopy to evaluate the rectal anastomosis for local recurrence for patients who have undergone an LAR. Anastomotic recurrence of rectal cancer has a much more favorable prognosis than local recurrence at other locations in the pelvis,^{141,142} although the optimal timing for surveillance of the rectal anastomosis is not known.

Following curative-intent surgery, post-treatment surveillance of patients with colorectal cancer is performed to evaluate for possible therapeutic complications, discover a recurrence that is potentially resectable for cure, and to identify new metachronous neoplasms at a preinvasive stage. Advantages of more intensive follow-up of Stage II and/or Stage III patients have been demonstrated prospectively in several studies²³⁶⁻²³⁸ and in 3 recent meta-analyses of randomized

controlled trials designed to compare low-intensity and high-intensity programs of surveillance.^{239,240-242} Other recent studies impacting on the issue of post-treatment surveillance of colorectal cancer include results from an analysis of data from 20,898 patients enrolled in 18 large adjuvant colon cancer randomized trials which demonstrated that 80% of recurrences were in the first 3 years after surgical resection of the primary tumor,²⁴³ and a population-based report indicating increased rates of resectability and survival in patients treated for local recurrence and distant metastases of colorectal cancer, thereby providing support for more intensive post-treatment follow-up in these patients.²⁴⁴ Nevertheless, controversies remain regarding selection of optimal strategies for following up patients after potentially curative colorectal cancer surgery.^{245,246}

The following panel recommendations for post-treatment surveillance pertain to patients with stage I-stage III disease who have undergone successful treatment (i.e. no known residual disease): history and physical examination every 3-6 months for 2 years, and then every 6 months for a total of 5 years; and a CEA test at baseline and every 3-6 months for 2 years,²⁴⁷ then every 6 months for the next 5 years for patients with disease staged as T2 or greater.^{242,247,248} Colonoscopy is recommended at approximately 1 year following resection (or at approximately 6 months post resection if not performed preoperatively due to obstructing lesion). Repeat colonoscopy is typically recommended at 3 years, and then every 5 years thereafter, unless follow-up colonoscopy indicates advanced adenoma (villous polyp, polyp > 1 cm or high grade dysplasia) in which case colonoscopy should be repeated in 1 year.²⁴⁹ More frequent colonoscopies may be indicated in patients who present with colorectal cancer before age 50. Proctoscopy should be considered every 6 months for 5 years to evaluate for local recurrence at the rectal anastomosis for patients who have undergone an LAR. Chest, abdominal and pelvic CT scans are recommended annually for the first 3 to 5 years in Stage II and III

patients.^{242,245} Routine PET scanning is not recommended and should not be obtained either as a routine pre-operative baseline study or for routine surveillance.

Initial follow-up office visits at 3 months intervals for history and physical examination may be more useful for patients diagnosed with Stage III disease, whereas patients with a diagnosis of Stage I disease may not need to be seen as frequently (i.e. can be seen once every 6 months). This principle also applies to CEA testing,²⁵⁰ which is used primarily to monitor for recurrence of the original disease (see section on Managing an Increasing CEA Level, below), although post-treatment CEA testing is recommended only if the patient is a potential candidate for further intervention.²⁴⁷ Surveillance colonoscopies are primarily aimed at identifying and removing metachronous polyps since data show that patients with a history of colorectal cancer have an increased risk of developing second cancers,²⁵¹ particularly in the first 2 years following resection. Furthermore, use of post-treatment surveillance colonoscopy has not been shown to improve survival through the early detection of recurrence of the original colorectal cancer.²⁴⁹ CT scan is recommended to monitor for the presence of potentially resectable metastatic lesions, primarily in the lung and the liver. Hence, CT scan is not routinely recommended in patients who are not candidates for potentially curative resection of liver or lung metastases.^{242,245} Post-treatment PET scan is not routinely recommended for surveillance of patients with resected early-stage colorectal cancer to detect recurrence of the original cancer.²⁴⁵ Furthermore, PET scan is not routinely recommended to detect metastatic disease in the absence of other evidence of such disease.

Managing an Increasing Carcinoembryonic Antigen Level

Managing patients with an elevated CEA level after resection should include colonoscopy, chest, abdominal, and pelvic CT scans, and consideration of a PET scan. If imaging study results are normal in the

face of a rising CEA, repeat CT scans are indicated every 3 months until either disease is identified or CEA level stabilizes or declines. The opinion of the panel regarding the usefulness of PET scan in the scenario of an elevated CEA with negative, good-quality CT scans was divided. Some favored use of PET in this scenario whereas others noted that the likelihood of PET identifying surgically curable disease in the setting of negative good-quality CT scans is vanishingly small. Use of PET scans in this scenario is permissible within these guidelines. The panel does not recommend the use of anti-CEA--radiolabeled scintigraphy.²⁵² In the event that surgically curable metastatic disease is identified on CT or MRI, the panel does recommend a PET scan before surgical resection to look for evidence of additional metastases that may change the status of patient resectability.

Summary

The NCCN Rectal Cancer Guidelines panel believes that a multidisciplinary approach, including representation from gastroenterology, medical oncology, surgical oncology, radiation oncology, and radiology is necessary for treating patients with rectal cancer. Adequate pathologic assessment of the resected lymph nodes is important with a goal of evaluating at least 12 nodes when possible. Patients with T1 or T2 lesions that are node-negative by endorectal ultrasound or endorectal or pelvic MRI and who meet carefully defined criteria can be managed with a transanal excision. A transabdominal resection is appropriate for all other rectal lesions. Preoperative chemoRT is preferred for the majority of patients with suspected or proven T3/T4 disease and/or regional node involvement and adjuvant chemotherapy is recommended. Patients with recurrent localized disease should be considered for resection with or without radiotherapy.

A patient with metastatic disease in the liver or lung should be considered for surgical resection if he or she is a candidate for surgery

and if complete resection (R0) or ablation can be achieved. Preoperative chemotherapy can be considered as initial therapy in patients with synchronous or metachronous resectable metastatic disease (ie, neoadjuvant therapy) or when a response to chemotherapy may convert a patient from an unresectable to resectable state (ie, conversion therapy). Another option for these patients is initial treatment with chemoRT. Resection should be followed by adjuvant therapy based on prior therapy received. The recommended post-treatment surveillance program for rectal cancer patients includes serial CEA determinations, as well as periodic chest, abdominal and pelvic CT scans, and periodic evaluations by colonoscopy and proctoscopy.

Recommendations for patients with previously untreated disseminated metastatic disease represent a continuum of care in which lines of treatment are blurred rather than discrete. Principles to consider at the start of therapy include pre-planned strategies for altering therapy for patients in both the presence and absence of disease progression, including plans for adjusting therapy for patients who experience certain toxicities. Recommended initial therapy for advanced or metastatic disease includes bevacizumab plus FOLFOX, FOLFIRI, capecitabine or 5-FU/LV. Chemotherapy options for patients with progressive disease are dependent on the choice of initial therapy and, for those patients able to tolerate intensive therapy, include FOLFIRI, CapeOX, FOLFOX or irinotecan alone or the combination of cetuximab with either irinotecan or FOLFIRI. Monotherapy with either cetuximab or panitumumab is also an option for patients not able to tolerate the combination of irinotecan plus cetuximab after first or second progression of disease. The panel endorses the concept that treating patients in a clinical trial has priority over standard or accepted therapy.

Discussion
Update in
Progress

References

1. Jemal A, Siegel R, Ward E, et al. Cancer statistics, 2008. *CA Cancer J Clin.* 2008;58:71-96.
2. Bonelli L, Martines H, Conio M, et al. Family history of colorectal cancer as a risk factor for benign and malignant tumours of the large bowel. A case-control study. *Int J Cancer.* 1988;41:513-517.
3. Ahsan H, Neugut AI, Garbowski GC, et al. Family history of colorectal adenomatous polyps and increased risk for colorectal cancer. *Ann Intern Med.* 1998;128:900-905.
4. Greene F, Page D, Fleming I, Fritz A. *AJCC Cancer Staging Manual.* New York: Springer-Verlag; 2002.
5. Compton CC, Greene FL. The staging of colorectal cancer: 2004 and beyond. *CA Cancer J Clin.* 2004;54:295-308.
6. Greene FL, Stewart AK, Norton HJ. A new TNM staging strategy for node-positive (stage III) colon cancer: an analysis of 50,042 patients. *Ann Surg.* 2002;236:416-421; discussion 421.
7. Compton CC. Updated protocol for the examination of specimens from patients with carcinomas of the colon and rectum, excluding carcinoid tumors, lymphomas, sarcomas, and tumors of the vermiform appendix: a basis for checklists. Cancer Committee. *Arch Pathol Lab Med.* 2000;124:1016-1025.
8. Compton CC. Key issues in reporting common cancer specimens: problems in pathologic staging of colon cancer. *Arch Pathol Lab Med.* Mar 2006;130:318-324.
9. Lai LL, Fuller CD, Kachnic LA, Thomas CR, Jr. Can pelvic radiotherapy be omitted in select patients with rectal cancer? *Semin Oncol.* 2006;33(6 Suppl 11):S70-74.
10. Nagtegaal ID, Marijnen CA, Kranenbarg EK, et al. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimeter but two millimeters is the limit. *Am J Surg Pathol.* 2002;26:350-357.
11. Wibe A, Rendedal PR, Svensson E, et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surg.* Mar 2002;89:327-334.
12. Stocchi L, Nelson H, Sargent DJ, et al. Impact of surgical and pathologic variables in rectal cancer: a United States community and cooperative group report. *J Clin Oncol.* 2001;19:3895-3902.
13. Glynne-Jones R, Mawdsley S, Novell JR. The clinical significance of the circumferential resection margin following preoperative pelvic chemo-radiotherapy in rectal cancer: why we need a common language. *Colorectal Dis.* 2006;8:800-807.
14. Adam IJ, Mohamdee MO, Martin IG, et al. Role of circumferential margin involvement in the local recurrence of rectal cancer. *Lancet.* 1994;344:707-711.
15. Mawdsley S, Glynne-Jones R, Grainger J, et al. Can histopathologic assessment of circumferential margin after preoperative pelvic chemoradiotherapy for T3-T4 rectal cancer predict for 3-year disease-free survival? *Int J Radiat Oncol Biol Phys.* 2005;63:745-752.
16. Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? *J Clin Oncol.* 2008;26:303-312.
17. Sarli L, Bader G, Iusco D, et al. Number of lymph nodes examined and prognosis of TNM stage II colorectal cancer. *Eur J Cancer.* 2005;41:272-279.
18. Newland RC, Dent OF, Lyttle MN, et al. Pathologic determinants of survival associated with colorectal cancer with lymph node metastases. A multivariate analysis of 579 patients. *Cancer.* 1994;73:2076-2082.
19. Wong SL, Hong J, Hollenbeck BK, et al. Hospital lymph node examination rates and survival after resection for colon cancer. *JAMA.* 2007;298:2149-2154.

20. Tepper JE, O'Connell MJ, Niedzwiecki D, et al. Impact of number of nodes retrieved on outcome in patients with rectal cancer. *J Clin Oncol*. 2001;19:157-163.
21. Pocard M, Panis Y, Malassagne B, et al. Assessing the effectiveness of mesorectal excision in rectal cancer: prognostic value of the number of lymph nodes found in resected specimens. *Dis Colon Rectum*. 1998;41:839-845.
22. Wichmann MW, Muller C, Meyer G, et al. Effect of preoperative radiochemotherapy on lymph node retrieval after resection of rectal cancer. *Arch Surg*. 2002;137:206-210.
23. Baxter NN, Morris AM, Rothenberger DA, Tepper JE. Impact of preoperative radiation for rectal cancer on subsequent lymph node evaluation: population based analysis. *Int J Rad Oncol Biol Phys*. 2005;61:426-431.
24. Meyers MO, Hollis DR, Mayer RJ, et al. Ratio of metastatic to examined lymph nodes is a powerful predictor of overall survival in rectal cancer: An analysis of Intergroup 0114. *J Clin Oncol*. 2007;25: No. 18S (June 20 suppl). Abstract 4008.
25. Turner RR, Nora DT, Trocha SD, Bilchik AJ. Colorectal carcinoma nodal staging. Frequency and nature of cytokeratin-positive cells in sentinel and nonsentinel lymph nodes. *Arch Pathol Lab Med*. 2003;127:673-679.
26. Wood TF, Nora DT, Morton DL, et al. One hundred consecutive cases of sentinel node mapping in early colorectal carcinoma. Detection of micrometastasis. *J Gastrointest Surg*. 2002;6:322-330.
27. Noura S, Yamamoto H, Miyake Y, et al. Immunohistochemical assessment of localization and frequency of micrometastases in lymph nodes of colorectal cancer. *Clin Cancer Res*. 2002;8:759-767.
28. Yasuda K, Adachi Y, Shiraishi N, et al. Pattern of lymph node micrometastasis and prognosis of patients with colorectal cancer. *Ann Surg Oncol*. 2001;8:300-304.
29. Braat AE, Oosterhuis JW, Moll FC, et al. Sentinel node detection after preoperative short-course radiotherapy in rectal carcinoma is not reliable. *Br J Surg*. 2005;92:1533-1538.
30. Cooper HS, Deppisch LM, Gourley WK, et al. Endoscopically removed malignant colorectal polyps: clinicopathologic correlations. *Gastroenterology*. 1995;108:1657-1665.
31. Nivatvongs S, Rojanasakul A, Reiman HM, et al. The risk of lymph node metastasis in colorectal polyps with invasive adenocarcinoma. *Dis Colon Rectum*. 1991;34:323-328.
32. Volk EE, Goldblum JR, Petras RE, et al. Management and outcome of patients with invasive carcinoma arising in colorectal polyps. *Gastroenterology*. 1995;109:1801-1807.
33. Ueno H, Mochizuki H, Hashiguchi Y, et al. Risk factors for an adverse outcome in early invasive colorectal carcinoma. *Gastroenterology*. 2004;127:385-394.
34. Seitz U, Bohnacker S, Seewald S, et al. Is endoscopic polypectomy an adequate therapy for malignant colorectal adenomas? Presentation of 114 patients and review of the literature. *Dis Colon Rectum*. 2004;47:1789-1796; discussion 1796-1787.
35. Garcia-Aguilar J, Hernandez de Anda E, et al. Endorectal ultrasound in the management of patients with malignant rectal polyps. *Dis Colon Rectum*. 2005;48:910-916; discussion 916-917.
36. Winawer SJ, Zauber AG, Fletcher RH, et al. Guidelines for colonoscopy surveillance after polypectomy: a consensus update by the US Multi-Society Task Force on Colorectal Cancer and the American Cancer Society. *CA Cancer J Clin*. 2006;56:143-159; quiz 184-145.
37. Nelson H, Petrelli N, Carlin A, et al. Guidelines 2000 for colon and rectal cancer surgery. *J Natl Cancer Inst*. 2001;93:583-596.
38. Kapiteijn E, Marijnen CA, Nagtegaal ID, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med*. 2001;345:638-646.

39. Baxter NN, Garcia-Aguilar J. Organ preservation for rectal cancer. *J Clin Oncol.* 2007;25:1014-1020.
40. Weiser MR, Landmann RG, Wong WD, et al. Surgical salvage of recurrent rectal cancer after transanal excision. *Dis Colon Rectum.* 2005;48:1169-1175.
41. Wiig JN, Larsen SG, Giercksky KE. Operative treatment of locally recurrent rectal cancer. *Recent Results Cancer Res.* 2005;165:136-147.
42. Bartram C, Brown G. Endorectal ultrasound and magnetic resonance imaging in rectal cancer staging. *Gastroenterol Clin North Am.* 2002;31:827-839.
43. Bipat S, Glas AS, Slors FJ, et al. Rectal cancer: local staging and assessment of lymph node involvement with endoluminal US, CT, and MR imaging--a meta-analysis. *Radiology.* 2004;232:773-783.
44. Klessen C, Rogalla P, Taupitz M. Local staging of rectal cancer: the current role of MRI. *Eur Radiol.* 2007;17:379-389.
45. Lahaye MJ, Engelen SM, Nelemans PJ, et al. Imaging for predicting the risk factors--the circumferential resection margin and nodal disease--of local recurrence in rectal cancer: a meta-analysis. *Semin Ultrasound CT MR.* 2005;26:259-268.
46. Beets-Tan RG, Vliegen RF, Beets GL. Magnetic resonance imaging of rectal cancer: what radiation oncologists need to know. *Front Radiat Ther Oncol.* 2004;38:1-12.
47. Guillem JG, Cohen AM. Current issues in colorectal cancer surgery. *Semin Oncol.* 1999;26:505-513.
48. You YN, Baxter NN, Stewart A, Nelson H. Is the increasing rate of local excision for stage I rectal cancer in the United States justified?: a nationwide cohort study from the National Cancer Database. *Ann Surg.* 2007;245:726-733.
49. Jayne DG, Guillou, Thorpe H, et al. Randomized trial of laparoscopic-assisted resection of colorectal carcinoma: 3-year results of the UK MRC CLASICC trial group. *J Clin Oncol.* 2007;25:3061-3068.
50. Ng SS, Leung KL, Lee JF, et al. Laparoscopic-assisted versus open abdominoperineal resection for low rectal cancer: a prospective randomized trial. *Ann Surg Oncol.* 2008;
51. Wagman LJ. Laparoscopic and open surgery for colorectal cancer: reaching equipoise? *J Clin Oncol.* 2007;25:2996-2998.
52. Heald RJ, Husband EM, Ryall RD. The mesorectum in rectal cancer surgery--the clue to pelvic recurrence? *Br J Surg.* 1982;69:613-616.
53. Marr R, Birbeck K, Garvican J, et al. The modern abdominoperineal excision: the next challenge after total mesorectal excision. *Ann Surg.* 2005;242:74-82.
54. Steup WH, Moriya Y and van de velde CJH. Patterns of lymphatic spread in rectal cancer. A topographical analysis on lymph node metastases. *Eur J Cancer.* 2002;38:911-918.
55. Schlag PM. Surgical sphincter preservation in rectal cancer. *Oncologist.* 1996;1:288-292.
56. Wagman R, Minsky BD, Cohen AM, et al. Sphincter preservation in rectal cancer with preoperative radiation therapy and coloanal anastomosis: long term follow-up. *Int J Radiat Oncol Biol Phys.* 1998;42:51-57.
57. Sauer R, Becker H, Hohenberger W, et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. *N Engl J Med.* 2004;351:1731-1740.
58. Kachnic LA. Should preoperative or postoperative therapy be administered in the management of rectal cancer? *Semin Oncol.* Dec 2006;33(6 Suppl 11):S64-69.
59. Madoff RD. Chemoradiotherapy for rectal cancer--when, why, and how? *N Engl J Med.* 2004;351:1790-1792.

60. Guillem JG, Diaz-Gonzalez JA, Minsky B, et al. cT3N0 rectal cancer: potential overtreatment with preoperative chemoradiotherapy is warranted. *J Clin Oncol.* 2008;26:368-373.
61. Improved survival with preoperative radiotherapy in resectable rectal cancer. Swedish Rectal Cancer Trial. *N Engl J Med.* 1997;336:980-987.
62. Colorectal Collaborative Group. Adjuvant radiotherapy for rectal cancer: a systematic overview of 8,507 patients from 22 randomised trials. *Lancet.* 2001;358:1291-1304.
63. Peeters KCMJ, Marijnen CAM, Nagtegaal ID, et al. The TME trial after a median follow-up of 6 years. *Ann Surg.* 2007;246:693-701.
64. Sebag-Montefiore D, Steele R, Quirke P, et al. Routine short course pre-op radiotherapy or selective post-op chemoradiotherapy for resectable rectal cancer? Preliminary results of the MRC CR07 randomised trial. *J Clin Oncol.* 2006;24:No. 18S (June 20 suppl). Abstract 3511.
65. Gerard JP, Conroy T, Bonnetain F, et al. Preoperative radiotherapy with or without concurrent fluorouracil and leucovorin in T3-4 rectal cancers: results of FFCD 9203. *J Clin Oncol.* 2006;24:4620-4625.
66. Glynne-Jones R, Sebag-Montefiore D. Role of neoadjuvant chemotherapy in rectal cancer: interpretation of the EXPERT study. *J Clin Oncol.* 2006;24:4664-4665; author reply 4665-4666.
67. Bosset JF, Calais G, Mineur L, et al. Enhanced tumorocidal effect of chemotherapy with preoperative radiotherapy for rectal cancer: preliminary results--EORTC 22921. *J Clin Oncol.* 2005;23:5620-5627.
68. Bosset JF, Collette L, Calais G, et al. Chemotherapy with preoperative radiotherapy in rectal cancer. *N Engl J Med.* 2006;355:1114-1123.
69. Collette L, Bosset JF, den Dulk M, et al. Patients with curative resection of cT3-4 rectal cancer after preoperative radiotherapy or radiochemotherapy: does anybody benefit from adjuvant fluorouracil-based chemotherapy? A trial of the EORTC Group. *J Clin Oncol.* 2007;25:4379-4386.
70. Bujko K, Kepka L, Michalski W, Nowacki MP. Does rectal cancer shrinkage induced by preoperative radio(chemo)therapy increase the likelihood of anterior resection? A systematic review of randomised trials. *Radiother Oncol.* 2006;80:4-12.
71. Wong RK, Tandan V, De Silva S, Figueredo A. Pre-operative radiotherapy and curative surgery for the management of localized rectal carcinoma. *Cochrane Database Syst Rev.* 2007(2):CD002102.
72. Peeters KC, van de Velde CJ, Leer JW, et al. Late side effects of short-course preoperative radiotherapy combined with total mesorectal excision for rectal cancer: increased bowel dysfunction in irradiated patients--a Dutch colorectal cancer group study. *J Clin Oncol.* 2005;23:6199-6206.
73. Tepper JE, O'Connell M, Niedzwiecki D, et al. Adjuvant therapy in rectal cancer: analysis of stage, sex, and local control--final report of intergroup 0114. *J Clin Oncol.* 2002;20:1744-1750.
74. Gunderson LL, Sargent DJ, Tepper JE, et al. Impact of T and N stage and treatment on survival and relapse in adjuvant rectal cancer: a pooled analysis. *J Clin Oncol.* 2004;22:1785-1796.
75. Johal BS, Phang P, McGahan C, et al. Treatment patterns and outcomes of patients with pT3N0 rectal cancer in a population-based setting. *J Clin Oncol.* 2007;25: No. 18S (June 20 suppl). Abstract 4039.
76. Smalley SR, Benedetti JK, Williamson SK, et al. Phase III trial of fluorouracil-based chemotherapy regimens plus radiotherapy in postoperative adjuvant rectal cancer: GI INT 0144. *J Clin Oncol.* 2006;24:3542-3547.
77. O'Connell MJ, Martenson JA, Wieand HS, et al. Improving adjuvant therapy for rectal cancer by combining protracted-infusion fluorouracil with radiation therapy after curative surgery. *N Engl J Med.* 1994;331:502-507.

78. Kim JS, Cho MJ, Song KS, Yoon WH. Preoperative chemoradiation using oral capecitabine in locally advanced rectal cancer. *Int J Radiat Oncol Biol Phys*. 2002;54:403-408.
79. Shi GG, Lin E, C. E, et al. Phase II study of capecitabine and radiotherapy plus concomitant boost in patients with locally advanced rectal cancer (LARC). *J Clin Oncol*. 2004;22:No. 14S (June 15 suppl). Abstract 3775.
80. Glynne-Jones R, Dunst J, Sebag-Montefiore D. The integration of oral capecitabine into chemoradiation regimens for locally advanced rectal cancer: how successful have we been? *Ann Oncol*. 2006;17:361-371.
81. De Paoli A, Chiara S, Luppi G, et al. A phase II study of capecitabine and pre-operative radiation therapy in resectable, locally advanced rectal cancer. *J Clin Oncol*. 2004;22:No. 14S (July 15 suppl). Abstract 3540.
82. Krishnan S, Janjan NA, Skibber JM, et al. Phase II study of capecitabine (Xeloda) and concomitant boost radiotherapy in patients with locally advanced rectal cancer. *Int J Radiat Oncol Biol Phys*. 2006;66:762-771.
83. Dupruis O, Vie B, Lledo G, et al. Preoperative treatment combining capecitabine with radiation therapy in rectal cancer : A GERCOR phase II study. *Oncology*. 2007;73:169-176.
84. Das P, Lin EH, Bhatia S, et al. Preoperative chemoradiotherapy with capecitabine versus protracted infusion 5-fluorouracil for rectal cancer: a matched pair analysis. *Int J Radiat Oncol Biol Phys*. 2006;66:1378-1383.
85. Andre T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med*. 2004;350:2343-2351.
86. Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med*. 2005;352:2696-2704.
87. Benson AB 3rd. New approaches to assessing and treating early-stage colon and rectal cancers: cooperative group strategies for assessing optimal approaches in early-stage disease. *Clin Cancer Res*. 2007;13:6913s-6920s.
88. Hong TS, Ritter MA, Tome WA, Harari PM. Intensity-modulated radiation therapy: emerging cancer treatment technology. *Br J Cancer*. 2005;92:1819-1824.
89. Meyer J, Czito B, Yin FF, Willett C. Advanced radiation therapy technologies in the treatment of rectal and anal cancer: intensity-modulated photon therapy and proton therapy. *Clin Colorectal Cancer*. 2007;6:348-356.
90. American College of Radiology. Practice Guideline for Intensity-modulated radiation therapy (IMRT). 2007; Available at: http://www.acr.org/SecondaryMainMenuCategories/quality_safety/guidelines/ro/imrt.aspx
91. Valenti V, Balducci M, Tororeto F, et al. Intraoperative radiotherapy: current thinking. *Eur J Surg Oncol*. 2002;28:180-185.
92. Hahnloser D, Haddock MG, Nelson H. Intraoperative radiotherapy in the multimodality approach to colorectal cancer. *Surg Oncol Clin N Am*. 2003;12:993-1013.
93. Willett CG, Czito BG, Tyler DS. Intraoperative radiation therapy. *J Clin Oncol*. 2007;25:971-977.
94. Moore HG, Gittleman AE, Minsky BD, et al. Rate of pathologic complete response with increased interval between preoperative combined modality therapy and rectal cancer resection. *Dis Colon Rectum*. 2004;47:279-286.
95. Habr-Gama A, Perez RO, Proscurshim I, et al. Interval between surgery and neoadjuvant chemoradiation therapy for distal rectal cancer: does delayed surgery have an impact on outcome? *Int J Radiat Oncol Biol Phys*. 2008;71:1181-1188.
96. Tulchinsky H, Shmuell E, Giger A, et al. An interval >7 weeks between neoadjuvant therapy and surgery improves pathologic

complete response and disease-free survival in patients with locally advanced rectal cancer. *Ann Surg Oncol*. 2008

97. Tran CL, Udani S, Holt A, Arnell T, et al. Evaluation of safety of increased time interval between chemoradiation and resection for rectal cancer. *Am J Surg*. 2006;192:873-877.

98. Wolmark N, Wieand HS, Hyams DM, et al. Randomized trial of postoperative adjuvant chemotherapy with or without radiotherapy for carcinoma of the rectum: National Surgical Adjuvant Breast and Bowel Project Protocol R-02. *J Natl Cancer Inst*. 2000;92:388-396.

99. Benson AB, 3rd. New approaches to the adjuvant therapy of colon cancer. *Oncologist*. 2006;11:973-980.

100. Sengupta S, Tjandra JJ. Local excision of rectal cancer: what is the evidence? *Dis Colon Rectum*. 2001;44:1345-1361.

101. Garcia-Aguilar J, Mellgren A, Sirivongs P, et al. Local excision of rectal cancer without adjuvant therapy: a word of caution. *Ann Surg*. 2000;231:345-351.

102. Willett CG, Compton CC, Shellito PC, Efird JT. Selection factors for local excision or abdominoperineal resection of early stage rectal cancer. *Cancer*. 1994;73:2716-2720.

103. Van Cutsem E, Nordlinger B, Adam R, et al. Towards a pan-European consensus on the treatment of patients with colorectal liver metastases. *Eur J Cancer*. 2006;42:2212-2221.

104. Yoo PS, Lopez-Soler RI, Longo WE, Cha CH. Liver resection for metastatic colorectal cancer in the age of neoadjuvant chemotherapy and bevacizumab. *Clin Colorectal Cancer*. 2006;6:202-207.

105. Kemeny N. Management of liver metastases from colorectal cancer. *Oncology (Williston Park)*. 2006;20:1161-1176, 1179; discussion 1179-1180, 1185-1166.

106. Muratore A, Zorzi D, Bouzari H, et al. Asymptomatic colorectal cancer with un-resectable liver metastases: immediate colorectal

resection or up-front systemic chemotherapy? *Ann Surg Oncol*. 2007;14:766-770.

107. Alberts SR, Horvath WL, Sternfeld WC, et al. Oxaliplatin, fluorouracil, and leucovorin for patients with unresectable liver-only metastases from colorectal cancer: a North Central Cancer Treatment Group phase II study. *J Clin Oncol*. 2005;23:9243-9249.

108. Fong Y, Cohen AM, Fortner JG, et al. Liver resection for colorectal metastases. *J Clin Oncol*. Mar 1997;15(3):938-946.

109. Tsai M, Su Y, Ho M, et al. Clinicopathological features and prognosis in resectable synchronous and metachronous colorectal liver metastases. *Ann Surg Oncol*. 2007;14:786-794.

110. Foster JH. Treatment of metastatic disease of the liver: a skeptic's view. *Semin Liver Dis*. 1984;4:170-179.

111. Stangl R, Altendorf-Hofmann A, et al. Factors influencing the natural history of colorectal liver metastases. *Lancet*. 1994;343:1405-1410.

112. Venook AP. The Kemeny article reviewed. *Oncology*. 2006;20:477-484.

113. Choti MA, Sitzmann JV, Tiburi MF, et al. Trends in long-term survival following liver resection for hepatic colorectal metastases. *Ann Surg*. 2002;235:759-766.

114. Pawlik TM, Poon RT, Abdalla EK, et al. Critical appraisal of the clinical and pathologic predictors of survival after resection of large hepatocellular carcinoma. *Arch Surg*. 2005;140:450-457; discussion 457-458.

115. Charnsangavej C, Clary B, Fong Y, et al. Selection of patients for resection of hepatic colorectal metastases: expert consensus statement. *Ann Surg Oncol*. 2006;13:1261-1268.

116. Vauthey JN, Zorzi D, Pawlik TM. Making unresectable hepatic colorectal metastases resectable--does it work? *Semin Oncol*. Dec 2005;32(6 Suppl 9):S118-122.

117. Pozzo C, Basso M, Cassano A, et al. Neoadjuvant treatment of unresectable liver disease with irinotecan and 5-fluorouracil plus folinic acid in colorectal cancer patients. *Ann Oncol.* 2004;15:933-939.
118. Altendorf-Hofmann A, Scheele J. A critical review of the major indicators of prognosis after resection of hepatic metastases from colorectal carcinoma. *Surg Oncol Clin N Am.* 2003;12:165-192.
- 119.. Pawlik TM, Schulick RD, Choti MA. Expanding criteria for resectability of colorectal liver metastases. *The Oncologist.* 2008;13:51-64.
120. Folprecht G, Grothey A, Alberts S, et al. Neoadjuvant treatment of unresectable colorectal liver metastases: correlation between tumour response and resection rates. *Ann Oncol.* 2005;16:1311-1319.
121. Abdalla EK, Vauthey JN, Ellis LM, et al. Recurrence and outcomes following hepatic resection, radiofrequency ablation, and combined resection/ablation for colorectal liver metastases. *Ann Surg.* 2004;239:818-825; discussion 825-817
122. Leonard GD, Brenner B, Kemeny NE. Neoadjuvant chemotherapy before liver resection for patients with unresectable liver metastases from colorectal carcinoma. *J Clin Oncol.* 2005;23:2038-2048.
123. Benoist S, Brouquet A, Penna C, et al. Complete response of colorectal liver metastases after chemotherapy: does it mean cure? *J Clin Oncol.* 2006;24:3939-3945.
124. Bilchik AJ, Poston G, Curley SA, et al. Neoadjuvant chemotherapy for metastatic colon cancer: a cautionary note. *J Clin Oncol.* 2005;23:9073-9078.
125. Pawlik TM, Scoggins CR, Zorzi D, et al. Effect of surgical margin status on survival and site of recurrence after hepatic resection for colorectal metastases. *Ann Surg.* 2005;241:715-722, discussion 722-714.
126. Elias D, Liberale G, Vernerey D, et al. Hepatic and extrahepatic colorectal metastases: when resectable, their localization does not matter, but their total number has a prognostic effect. *Ann Surg Oncol.* 2005;12:900-909.
127. Fong Y, Salo J. Surgical therapy of hepatic colorectal metastasis. *Semin Oncol.* 1999;26:514-523.
128. Adam R, Delvart V, Pascal G, et al. Rescue surgery for unresectable colorectal liver metastases downstaged by chemotherapy: a model to predict long-term survival. *Ann Surg.* 2004;240:644-657; discussion 657-648.
129. Delaunoy T, Alberts SR, Sargent DJ, et al. Chemotherapy permits resection of metastatic colorectal cancer: experience from Intergroup N9741. *Ann Oncol.* 2005;16:425-429.
130. Scappaticci FA, Fehrenbacher L, Cartwright T, et al. Surgical wound healing complications in metastatic colorectal cancer patients treated with bevacizumab. *J Surg Oncol.* 2005;91:173-180.
131. Package Insert. Bevacizumab. South San Francisco, CA, Genentech, Inc. October 2006.
132. Gruenberger B, Tamandl D, Schueller J, et al. Bevacizumab, capecitabine, and oxaliplatin as neoadjuvant therapy for patients with potentially curable metastatic colorectal cancer. *J Clin Oncol.* 2008;26:1830-1835.
133. Reddy SK, Morse MA, Hurwitz HI, et al. Addition of bevacizumab to irinotecan- and oxaliplatin-based preoperative chemotherapy regimens does not increase morbidity after resection of colorectal liver metastases. *J Am Coll Surg.* 2008;206:96-106.
134. Lee WS, Yun SH, Chun HK, et al. Pulmonary resection for metastases from colorectal cancer: prognostic factors and survival. *Int J Colorectal Dis.* Nov 16 2006.
135. Headrick JR, Miller DL, Nagorney DM, et al. Surgical treatment of hepatic and pulmonary metastases from colon cancer. *Ann Thorac Surg.* 2001;71:975-979; discussion 979-980.

136. Kemeny N. Management of liver metastases from colorectal cancer. *Oncology (Williston Park)*. 2006;20:1161-1176, 1179; discussion 1179-1180, 1185-1166.
137. Kemeny N, Huang Y, Cohen AM, et al. Hepatic arterial infusion of chemotherapy after resection of hepatic metastases from colorectal cancer. *N Engl J Med*. 1999;341:2039-2048.
138. Kemeny NE, Gonen M. Hepatic arterial infusion after liver resection. *N Engl J Med*. 2005;352:7:734-735.
139. Nordlinger B, Sorbye H, Glimelius B, et al. Perioperative chemotherapy with FOLFOX4 and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC Intergroup trial 40983): a randomized controlled trial. *Lancet*. 2008;37:1007-1016.
140. Yu T-K, Bhosale PR, Crane CH, et al. Patterns of locoregional recurrence after surgery and radiotherapy or chemoradiation for rectal cancer. *Int J Rad Oncol Biol Phys*. 2008;
141. Lowy AM, Rich TA, Skibber JM, et al. Preoperative infusional chemoradiation, selective intraoperative radiation, and resection for locally advanced pelvic recurrence of colorectal adenocarcinoma. *Ann Surg*. 1996;223:177-185.
142. Hoffman JP, Riley L, Carp NZ, Litwin S. Isolated locally recurrent rectal cancer: a review of incidence, presentation, and management. *Semin Oncol*. 1993;20:506-519.
143. Pelosi E, Deandreis D. The role of 18F-fluoro-deoxy-glucose positron emission tomography (FDG-PET) in the management of patients with colorectal cancer. *Eur J Surg Oncol*. 2007;33:1-6.
144. Mohiuddin M, Marks G, Marks J. Long-term results of reirradiation for patients with recurrent rectal carcinoma. *Cancer*. 2002;95:1144-1150.
145. Valentini V, Morganti AG, Gambacorta MA, et al. Preoperative hyperfractionated chemoradiation for locally recurrent rectal cancer in patients previously irradiated to the pelvis: A multicentric phase II study. *Int J Radiat Oncol Biol Phys*. 2006; 64:1129-1139.
146. Das P, Delclos ME, Skibber JM, et al. Hyperfractionated accelerated radiotherapy for rectal cancer in patients with prior pelvic irradiation. *Int J Rad Oncol Biol Phys*. 2007; 69:S277.
147. Yan TD, Black D, Savady R, Sugarbaker PH. Systemic review on the efficacy of cytoreductive surgery combined with perioperative intraperitoneal chemotherapy for peritoneal carcinomatosis from colorectal carcinoma. *J Clin Oncol*. 2006;24:4011-4019.
148. Esquivel J, Sticca R, Sugarbaker P, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy in the management of peritoneal surface malignancies of colonic origin: a consensus statement. Society of Surgical Oncology. *Ann Surg Oncol*. 2007;14:128-133.
149. Kelly H, Goldberg RM. Systemic therapy for metastatic colorectal cancer: current options, current evidence. *J Clin Oncol*. 2005;23:4553-4560.
150. Goldberg RM. Therapy for metastatic colorectal cancer. *Oncologist*. 2006;11:981-987.
151. Goldberg RM, Rothenberg ML, Van Cutsem E, et al. The continuum of care: a paradigm for the management of metastatic colorectal cancer. *Oncologist*. 2007;12:38-50.
152. Goldberg RM, Sargent DJ, Morton RF, et al. Randomized controlled trial of reduced-dose bolus fluorouracil plus leucovorin and irinotecan or infused fluorouracil plus leucovorin and oxaliplatin in patients with previously untreated metastatic colorectal cancer: a North American Intergroup Trial. *J Clin Oncol*. 2006;24:3347-3353.
153. Douillard JY, Cunningham D, Roth AD, et al. Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: a multicentre randomised trial. *Lancet*. 2000;355:1041-1047.

154. Fuchs CS, Moore MR, Harker G, et al. Phase III comparison of two irinotecan dosing regimens in second-line therapy of metastatic colorectal cancer. *J Clin Oncol.* 2003;21:807-814.
155. Petrelli N, Herrera L, Rustum Y, et al. A prospective randomized trial of 5-fluorouracil versus 5-fluorouracil and high-dose leucovorin versus 5-fluorouracil and methotrexate in previously untreated patients with advanced colorectal carcinoma. *J Clin Oncol.* 1987;5:1559-1565.
156. Van Cutsem E, Twelves C, Cassidy J, et al. Oral capecitabine compared with intravenous fluorouracil plus leucovorin in patients with metastatic colorectal cancer: results of a large phase III study. *J Clin Oncol.* 2001;19:4097-4106.
157. Van Cutsem E, Hoff PM, Harper P, et al. Oral capecitabine vs intravenous 5-fluorouracil and leucovorin: integrated efficacy data and novel analyses from two large, randomised, phase III trials. *Br J Cancer.* 2004;90:1190-1197.
158. Buroker TR, O'Connell MJ, Wieand HS, et al. Randomized comparison of two schedules of fluorouracil and leucovorin in the treatment of advanced colorectal cancer. *J Clin Oncol.* 1994;12:14-20.
159. de Gramont A, Bosset JF, Milan C, et al. Randomized trial comparing monthly low-dose leucovorin and fluorouracil bolus with bimonthly high-dose leucovorin and fluorouracil bolus plus continuous infusion for advanced colorectal cancer: a French intergroup study. *J Clin Oncol.* 1997;15:808-815.
160. Maindrault-Goebel F, Louvet C, Andre T, et al. Oxaliplatin added to the simplified bimonthly leucovorin and 5-fluorouracil regimen as second-line therapy for metastatic colorectal cancer (FOLFOX6). GERCOR. *Eur J Cancer.* 1999;35:1338-1342.
161. Van Cutsem E, Peeters M, Siena S, et al. Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol.* 2007; 25:1658-1664...
162. Cunningham D, Humblet Y, Siena S, et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med.* 2004;351:337-345.
163. Bartlett DL, Berlin J, Lauwers GY, et al. Chemotherapy and regional therapy of hepatic colorectal metastases: expert consensus statement. *Ann Surg Oncol.* 2006;13:1284-1292.
164. van Cutsem E. Challenges in the use of epidermal growth factor receptor inhibitors in colorectal cancer. *Oncologist.* 2006;11:1010-1017.
165. Raymond E, Faivre S, Woynarowski JM, Chaney SG. Oxaliplatin: mechanism of action and antineoplastic activity. *Semin Oncol.* 1998;25(Suppl 5):4-12.
166. O'Dwyer PJ. The present and future of angiogenesis-directed treatments of colorectal cancer. *Oncologist.* 2006;11:992-998.
167. Lentz F, Tran A, Rey E, et al. Pharmacogenomics of fluorouracil, irinotecan, and oxaliplatin in hepatic metastases of colorectal cancer: clinical implications. *Am J Pharmacogenomics.* 2005;5:21-33.
168. Rothenberg ML, Blanke CD. Topoisomerase I inhibitors in the treatment of colorectal cancer. *Semin Oncol.* 1999;26:632-639.
169. Goldberg RM, Sargent DJ, Morton RF, et al. A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol.* 2004;22:23-30.
170. Tournigand C, Andre T, Achille E, et al. FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: a randomized GERCOR study. *J Clin Oncol.* 2004;22:229-237.
171. Delaunoy T, Goldberg RM, Sargent DJ, et al. Mortality associated with daily bolus 5-fluorouracil/leucovorin administered in combination with either irinotecan or oxaliplatin: results from Intergroup Trial N9741. *Cancer.* 2004;101:2170-2176.

172. Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Br J Cancer*. 2002;87:393-399.
173. de Gramont A, Figer A, Seymour M, et al. Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol*. 2000;18:2938-2947.
174. Colucci G, Gebbia V, Paoletti G, et al. Phase III randomized trial of FOLFIRI versus FOLFOX4 in the treatment of advanced colorectal cancer: a multicenter study of the Gruppo Oncologico Dell'Italia Meridionale. *J Clin Oncol*. 2005;23:4866-4875.
175. Cassidy J, Clarke S, Rubio D, et al. First efficacy and safety results from XELOX-1/NO16966, a randomized 2x2 factorial phase III trial of XELOX vs. FOLFOX6 + bevacizumab or placebo in first-line metastatic colorectal cancer (MCR). *Ann Oncol*. 2006;17:(suppl 9). LBA 3.
176. Cassidy J, Tabernero J, Twelves C, et al. XELOX (capecitabine plus oxaliplatin): active first-line therapy for patients with metastatic colorectal cancer. *J Clin Oncol*. 2004;22:2084-2091.
177. Arkenau H, Schmoll H, Kubicka S, et al. Infusional 5-fluorouracil/folinic acid plus oxaliplatin (FUFOX) versus capecitabine plus oxaliplatin (CAPOX) as first line treatment of metastatic colorectal cancer (MCR): Results of the safety and efficacy analysis. *J Clin Oncol*. 2005; 23: 16S (June 1 suppl). Abstract 3507.
178. Andre T, Louvet C, Maindrault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *GERCOR*. *Eur J Cancer*. 1999;35:1343-1347.
179. Jager E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. Study Group for Palliative Treatment of Metastatic Colorectal Cancer Study Protocol 1. *J Clin Oncol*. 1996;14:2274-2279.
180. Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from National Surgical Adjuvant Breast and Bowel Project protocol C-03. *J Clin Oncol*. 1993;11:1879-1887.
181. Kabbinavar FF, Schulz J, McCleod M, et al. Addition of bevacizumab to bolus fluorouracil and leucovorin in first-line metastatic colorectal cancer: results of a randomized phase II trial. *J Clin Oncol*. 2005;23:3697-3705.
182. Hurwitz HI, Fehrenbacher L, Hainsworth JD, et al. Bevacizumab in combination with fluorouracil and leucovorin: an active regimen for first-line metastatic colorectal cancer. *J Clin Oncol*. 2005;23:3502-3508.
183. Hochster H, Hart LL, Ramanathan R, et al. Safety and efficacy of oxaliplatin/fluoropyrimidine regimens with or without bevacizumab as first-line treatment of metastatic colorectal cancer (mCRC): Final analysis of the TREE study. *J Clin Oncol*. 2006;24: N0. 18S (June 20 suppl). Abstract 3510.
184. Hurwitz H, Fehrenbacher L, Novotny W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med*. 2004;350:2335-2342.
185. Kabbinavar FF, Hambleton J, Mass RD, et al. Combined analysis of efficacy: the addition of bevacizumab to fluorouracil/leucovorin improves survival for patients with metastatic colorectal cancer. *J Clin Oncol*. 2005;23:3706-3712.
186. Saltz LB, Clarke S, Diaz-Rubio E, et al. Bevacizumab in combination with XELOX or FOLFOX4: Efficacy results from XELOX-1/NO16966, a randomized phase III trial in the first-line treatment of metastatic colorectal cancer. Paper presented at: Gastrointestinal Cancers Symposium 2007. Abstract 238.
187. Cassidy J, Clarke S, Diaz-Rubio E, et al. XELOX vs. FOLFOX4: Update of efficacy results from XELOX-1/NO16966, a randomized phase III trial of first-line treatment for patients with metastatic colorectal cancer. Paper presented at: Gastrointestinal Cancers Symposium 2008. Abstract 341.

188. Saltz LB, Clarke S, Díaz-Rubio E, et al. Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in metastatic colorectal cancer: a randomized phase III study. *J Clin Oncol*. 2008;26:2013-2039.
189. Giantonio BJ, Meropol NJ, Catalano PJ, et al. Magnitude of progression-free survival improvement and treatment duration in metastatic colorectal cancer for bevacizumab in combination with oxaliplatin-containing regimens: an analysis for two phase II studies. *J Clin Oncol*. 2007;25:No. 18S (June 20 suppl). Abstract 4073.
190. Fuchs CS, Marshall J, Mitchell EP, et al. Updated results of BICC-C study comparing first-line irinotecan /fluoropyrimidine combinations +/- celecoxib in mCRC: Clinical data cut-off September 1, 2006. Paper presented at: Gastrointestinal Cancers Symposium, 2007; Abstract 276.
191. Fuchs CJ, Marshall J, Barrueco J, et al. Randomized, controlled trial of irinotecan plus infusional, bolus, or oral fluoropyrimidines in first-line treatment of metastatic colorectal cancer: updated results from the BICC-C study. *J Clin Oncol*. 2008;25:689-690.
192. Sobero A, Ackland S, Carrion R, et al. Efficacy and safety of bevacizumab in combination with irinotecan and infusional 5-FU as first-line treatment for patients with metastatic colorectal cancer. *J Clin Oncol*. 2006;24:No. 18S (June 20 suppl). Abstract 3544.
193. Giantonio B, Catalano P, Meropol N, et al. Bevacizumab in combination with oxaliplatin, fluorouracil, and leucovorin (FOLFOX4) for previously treated metastatic colorectal cancer: results from the Eastern Cooperative Oncology Group Study E3200. *J Clin Oncol*. 2007;25:1539-1544.
194. Package Insert. Bevacizumab (Avastin®). South San Francisco, CA, Genentech, Inc. October 2006.
195. Scappaticci FA, Fehrenbacher L, Cartwright T, et al. Surgical wound healing complications in metastatic colorectal cancer patients treated with bevacizumab. *J Surg Oncol*. 2005;91:173-180.
196. Cannistra SA, Matulonis UA, Penson RT, et al. Phase II study of bevacizumab in patients with platinum-resistant ovarian cancer or peritoneal serous cancer. *J Clin Oncol*. 2007;25:5150-5152.
197. Package insert. Capecitabine (Xeloda®), Nutley, NJ, Roche Laboratories Inc., June 2005.
198. Haller D, Cassidy J, Clarke S, et al. Tolerability of fluoropyrimidines appears to differ by region. *J Clin Oncol*. 2006;24:16S (June 20 suppl). Abstract 3514.
199. Schmoll HJ, Arnold D. Update on capecitabine in colorectal cancer. *Oncologist*. 2006;11:1003-1009.
200. Package Insert. Irinotecan hydrochloride injection (Camptosar®), New York, NY, Pfizer, June 2006.
201. Innocenti F, Undevia SD, Iyer L, et al. Genetic variants in the UDP-glucuronosyltransferase 1A1 gene predict the risk of severe neutropenia of irinotecan. *J Clin Oncol*. 2004;22:1382-1388.
202. LabCorp Capsule. UGT1A1 irinotecan toxicity. Managing medication dosing and predicting response to treatment of cancer with irinotecan (Camptosar, CPT-11). 2006:Available at www.lapcorp.com.
203. O'Dwyer PJ, Catalano RB. Uridine diphosphate glucuronosyltransferase (UGT) 1A1 and irinotecan: practical pharmacogenomics arrives in cancer therapy. *J Clin Oncol*. 2006;24:4534-4538.
204. Package insert. Oxaliplatin (Eloxatin®), Bedford, OH. Ben Venue Laboratories, November 2004.
205. Tournigand C, Cervantes A, Figer A, et al. OPTIMOX1: a randomized study of FOLFOX4 or FOLFOX7 with oxaliplatin in a stop-and-Go fashion in advanced colorectal cancer--a GERCOR study. *J Clin Oncol*. 2006;24:394-400.
206. Maindrault-Goebel F, Lledo G, Ghibaudel B, et al. Final results for OPTIMOX-2, a large randomized phase II study of maintenance therapy or chemotherapy-free intervals after FOLFOX in patients with

metastatic colorectal cancer: a GERCOR study. *J Clin Oncol.* 2007;25: No. 18S. Abstract 4013.

207. Grothey A, Sargent D, Goldberg RM, Schmol HJ. Survival of patients with advanced colorectal cancer improves with the availability of fluorouracil-leucovorin, irinotecan, and oxaliplatin in the course of treatment. *J Clin Oncol.* 2004;22:1209-1214.

208. Cunningham D, Pyrhonen S, James RD, et al. Randomised trial of irinotecan plus supportive care versus supportive care alone after fluorouracil failure for patients with metastatic colorectal cancer. *Lancet.* 1998;352:1413-1418.

209. Rougier P, Van Cutsem E, Bajetta E, et al. Randomised trial of irinotecan versus fluorouracil by continuous infusion after fluorouracil failure in patients with metastatic colorectal cancer. *Lancet.* 1998;352:1407-1412.

210. Pitot H, Rowland K, DJ S, et al. N9841: A randomized phase III equivalence trial of irinotecan (CPT-11) versus oxaliplatin/5-fluorouracil (5FU)/leucovorin (FOLFOX4) in patients (pts) with advanced colorectal cancer (CRC) previously treated with 5FU. *J Clin Oncol.* 2005;23:No. 16S (June 1 suppl). Abstract 3506.

211. Baselga J, Rosen N. Determinants of RASistance to anti-epidermal growth factor receptor agents. *J Clin Oncol.* 2008;26:1582-1584.

212. Amado RG, Wolf M, Peeters M, et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol.* 2008;26:1626-1634.

213. Khambata-Ford S, Garrett CR, Meropol NJ, et al. Expression of epiregulin and amphiregulin and K-ras mutation status predict disease control in metastatic colorectal cancer patients treated with cetuximab. *J Clin Oncol.* 2007;25:3230-3237.

214. De Roock W, Piessevaux H, De Schutter J, et al. KRAS wild-type state predicts survival and is associated to early radiological response

in metastatic colorectal cancer treated with cetuximab. *Ann Oncol.* 2008;19:508-515.

215. Punt CJ, Tol J, Rodenburg CJ, et al. Randomized phase III study of capecitabine, oxaliplatin, and bevacizumab with or without cetuximab in advanced colorectal cancer, the CAIRO2 study of the Dutch Colorectal Cancer Group. *J Clin Oncol.* 2008;26 (May 20 suppl): Abstract LBA 4011.

216. Bokemeyer C, Bondarenko I, Hartmann JT, et al. KRAS status and efficacy of first-line treatment of patients with metastatic colorectal cancer with FOLFOX with or without cetuximab. The OPUS Experience. *J Clin Oncol.* 2008;26 (May 20 suppl): Abstract 4000.

217. Tejpar S, Peeters M, Humblet Y, et al. Relationship of efficacy with KRAS status (wild type versus mutant) in patients with irinotecan-refractory metastatic colorectal cancer, treated with irinotecan and escalating doses of cetuximab: The EVEREST experience (preliminary data). *J Clin Oncol.* 2008;26 (May 20 suppl): Abstract 4001.

218. Van Cutsem E, Lang I, Dhaens G, et al. KRAS status and efficacy in the first-line treatment of patients with metastatic colorectal cancer treated with FOLFIRI with or without cetuximab: The CRYSTAL experience. *J Clin Oncol.* 2008;26 (May 20 suppl): Abstract 2

219. Lièvre A, Bachet JB, Boige V, et al. KRAS mutations as an independent prognostic factor in patients with advanced colorectal cancer treated with cetuximab. *J Clin Oncol.* 2008;26:374-379.

220. Karapetis CS, Khambata-Ford S, Jonker DJ, et al. K-ras mutations and benefit from cetuximab in advanced colorectal cancer. *N Engl J Med.* 2008;359:1757-1765.

221. Artale S, Sartore-Bianchi A, Veronese SM, et al. Mutations in KRAS and BRAF in primary and matched metastatic sites of colorectal cancer. *J Clin Oncol.* 2008;26:4217-4219.

222. Saltz LB, Meropol NJ, Loehrer PJ, Sr., et al. Phase II trial of cetuximab in patients with refractory colorectal cancer that expresses the epidermal growth factor receptor. *J Clin Oncol.* 2004;22:1201-1208.

223. Saltz L, Rubin M, Hochster H, et al. Cetuximab (IMC-C225) plus irinotecan (CPT-11) is active in CPT-11-refractory colorectal cancer (CRC) that expresses epidermal growth factor receptor (EGFR). *Proc Am Soc Clin Oncol*. 2001; 20:Abstract 7.
224. Jonker DJ, O'Callaghan CJ, Karapetis CS, et al. Cetuximab for the treatment of colorectal cancer. *J Clin Oncol*. 2007;25:2040-2047.
225. Sobrero AF, Maurel J, Fehrenbacher L, et al. EPIC: Phase III trial of cetuximab plus irinotecan after fluoropyrimidine and oxaliplatin failure in patients with metastatic colorectal cancer. *J Clin Oncol*. 2008;26:2311-2319.
226. Hecht JR, Mitchell T, Chidiac C, et al. An updated analysis of safety and efficacy of oxaliplatin/bevacizumab +/- panitumumab for first-line treatment of metastatic colorectal cancer from a randomized, controlled trial (PACCE). *J Clin Oncol*. 2008 Gastrointestinal Cancers Symposium. Abstract 273.
227. Package Insert. Cetuximab (Erbix[®]), Branchburg, NJ, Imclone Systems, Inc. 2004.
228. Package Insert. Panitumumab (Vectibix[™]), Thousand Oaks, CA, Amgen, September 2006.
229. Helbling D, Borner M. Successful challenge with the fully human EGFR antibody panitumumab following an infusion reaction with the chimeric EGFR antibody cetuximab. *Ann Oncol*. 2007;18:963-964.
230. Heun J, Holen K. Treatment with panitumumab after a severe infusion reaction to cetuximab in a patient with metastatic colorectal cancer: a case report. *Clin Colorectal Cancer*. 2007;6:529-531.
231. Berlin J, van Cutsem E, Peeters M, et al. Predictive value of skin toxicity severity for response to panitumumab in patients with metastatic colorectal cancer: a pooled analysis of five clinical trials. *J Clin Oncol*. 2007;25: No. 18S (June 20 suppl). Abstract 4134.
232. van Cutsem E, Nowacki M, Lang I, et al. Randomized phase III study of irinotecan and 5-FU/FA with or without cetuximab in the first-line treatment of patients with metastatic colorectal cancer: the CRYSTAL trial. 2007; *J Clin Oncol*. 2007;25: No. 18S (June 20 suppl). Abstract 4000.
233. Hecht J, Mitchell E, Baranda J, et al. Panitumumab antitumor activity in patients (pts) with metastatic colorectal cancer (mCRC) expressing low (1-9%) or negative (<1%) levels of epidermal growth receptor (EGFr). *J Clin Oncol*. 2006;24: No. 18S (June 1 suppl). Abstract 3506.
234. Goldberg RM, Kohne GH, Seymour MT, et al. A pooled safety and efficacy analysis examining the effect of performance status on outcomes in nine first-line treatment trials of 6,286 patients with metastatic colorectal cancer. *J Clin Oncol*. 2007;25: No. 18S (June 20 suppl). Abstract 4011.
235. Hoff PM, Pazdur R, Lassere Y, et al. Phase II study of capecitabine in patients with fluorouracil-resistant metastatic colorectal carcinoma. *J Clin Oncol*. 2004;22:2078-2083.
236. Pietra N, Sarli L, Costi R, et al. Role of follow-up in management of local recurrences of colorectal cancer: a prospective, randomized study. *Dis Colon Rectum*. 1998;41:1127-1133.
237. Secco GB, Fardelli R, Gianquinto D, et al. Efficacy and cost of risk-adapted follow-up in patients after colorectal cancer surgery: a prospective, randomized and controlled trial. *Eur J Surg Oncol*. 2002;28:418-423.
238. Rodriguez-Moranta F, Salo J, Arcusa A, et al. Postoperative surveillance in patients with colorectal cancer who have undergone curative resection: a prospective, multicenter, randomized, controlled trial. *J Clin Oncol*. 2006;24:386-393.
239. Figueredo A, Charette ML, Maroun J, et al. Adjuvant therapy for stage II colon cancer: a systematic review from the Cancer Care Ontario Program in evidence-based care's gastrointestinal cancer disease site group. *J Clin Oncol*. 2004;22:3395-3407.
240. Renehan AG, Egger M, Saunders MP, O'Dwyer ST. Impact on survival of intensive follow up after curative resection for colorectal

cancer: systematic review and meta-analysis of randomised trials. *BMJ*. 2002;324:813.

241. Jeffery M, Hickey B, Hider P. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database Syst Rev*. 2007(1):CD002200.

242. Desch CE, Benson AB, 3rd, Somerfield MR, et al. Colorectal cancer surveillance: 2005 update of an American Society of Clinical Oncology practice guideline. *J Clin Oncol*. 2005;23:8512-8519.

243. Sargent DJ, Wieand HS, Haller DG, et al. Disease-free survival versus overall survival as a primary end point for adjuvant colon cancer studies: individual patient data from 20,898 patients on 18 randomized trials. *J Clin Oncol*. 2005;23:8664-8670.

244. Guyot F, Faivre J, Manfredi S, et al. Time trends in the treatment and survival of recurrences from colorectal cancer. *Ann Oncol*. 2005;16:756-761.

245. Pfister DG, Benson AB, 3rd, Somerfield MR. Clinical practice. Surveillance strategies after curative treatment of colorectal cancer. *N Engl J Med*. 2004;350:2375-2382.

246. Li Destri G, Di Cataldo A, Puleo S. Colorectal cancer follow-up: useful or useless? *Surg Oncol*. 2006;15:1-12.

247. Locker GY, Hamilton S, Harris J, et al. ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal cancer. *J Clin Oncol*. 2006;24:5313-5327.

248. Macdonald JS. Carcinoembryonic antigen screening: pros and cons. *Semin Oncol*. 1999;26:556-560.

249. Rex DK, Kahi CJ, Levin B, et al. Guidelines for colonoscopy surveillance after cancer resection: a consensus update by the American Cancer Society and US Multi-Society Task Force on Colorectal Cancer. *CA Cancer J Clin*. 2006;56:160-167; quiz 185-166.

250. Grossmann I, de Bock GH, Meershoek-Klein Kranenbarg WM, et al. Carcinoembryonic antigen (CEA) measurement during follow-up for

rectal carcinoma is useful even if normal levels exist before surgery. A retrospective study of CEA values in the TME trial. *Eur J Surg Oncol*. 2007;33:183-187.

251. Green RJ, Metlay JP, Propert K, et al. Surveillance for second primary colorectal cancer after adjuvant chemotherapy: an analysis of Intergroup 0089. *Ann Intern Med*. 2002;136:261-269.

252. Moffat FL, Jr., Pinsky CM, Hammershaimb L, et al. Clinical utility of external immunoscintigraphy with the IMMU-4 technetium-99m Fab' antibody fragment in patients undergoing surgery for carcinoma of the colon and rectum: results of a pivotal, phase III trial. The Immunomedics Study Group. *J Clin Oncol*. 1996;14:2295-2305.

Discussion
update in
progress